Neomycin Sulfate and Oxytetracycline Sheep Ration B Type C Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS FOR USE

For treatment of bacterial enteritis caused by <u>Escherichia coli</u> and bacterial pneumonia caused by Pasteurella multocida susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by Escherichia coli susceptible to neomycin.

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate		grams/ton.*
Oxytetracycline (from oxyt	tetracycline dihydrate base) equivalent to	grams oxytetracycline hydrochloride/ton.3

GUARANTEED ANALYSIS

Crude Protein, (Min)XX.XX%
NPN, (Max) ¹ X.XX%
Crude Fat, (Min) X.XX%
Crude Fiber, (Max) X.XX%
Calcium, (Min) X.XX%
Calcium, (Max) X.XX%
Phosphorus, (Min) X.XX%
Salt, (Min) ¹ X.XX%
Salt, (Max) ¹ X.XX%
Sodium, (Min) ² X.XX%
Sodium, (Max) ² X.XX%
Copper, (Min) ³ ppm
Copper, (Max) ³ ppm
Selenium, (Min) ⁴ ppm
Vitamin A, (Min) ⁴ IU/LB

¹If added


INGREDIENTS


Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed at 10 mg/lb body weight/day continuously for 7 - 14 days as the sole ration. Treatment should continue 24 to 48 hours beyond remission of disease symptoms. For example, the table below with the stated drug concentration, body weight and feed consumption will provide sheep with 10 mg/lb body weight/ day dosage.

Concentration of Oxytetracycline and Neomycin Sulfate (g/ton) in Type C Feed	Body Weight (lb)	Type C Feed Consumption (lb)
400	60	3.00
500	80	3.20
600	100	3.33





*The final printed medicated feed label must state a single drug concentration.

Net Weight____lb. (_____ kg.)

Manufactured by

Blue Bird Feed Company

Blue Bird, MD 00000

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

³If added, or if total copper exceeds 20 ppm

⁴Other than precursor of Vitamin A, if added