

**Neomycin Sulfate and Oxytetracycline Turkey Ration D
Type C Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS FOR USE

For control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate..... grams/ton.*
Oxytetracycline (from oxytetracycline dihydrate base) equivalent to ____ grams oxytetracycline hydrochloride/ton.*

GUARANTEED ANALYSIS

Crude Protein, (Min).....	XX.XX%
Lysine, (Min).....	X.XX%
Methionine, (Min).....	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min).....	X.XX%
Calcium, (Max).....	X.XX%
Phosphorus, (Min).....	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed at 25 mg/lb body weight/day continuously for 7-14 days as the sole ration. For example, the table below with the stated drug concentration, body weight and feed consumption will provide turkeys with 25 mg/lb body weight/day dosage.

Concentration of Oxytetracycline and Neomycin Sulfate (g/ton) in Type C Feed	Body Weight (lb)	Type C Feed Consumption (lb)
800	10	.63
900	15	0.83
1000	20	1.00

RESIDUE WARNING: Do not feed to turkeys producing eggs for human consumption. Withdraw 5 days before slaughter.

*The final printed medicated feed label must state a single drug concentration.

Net Weight _____ lb. (_____ kg.)

Manufactured by
Blue Bird Feed Company
Blue bird, MD 00000