

**Chlortetracycline Cattle Supplement
Type B Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated cattle feed.

ACTIVE INGREDIENT

Chlortetracycline.....2-40 g/lb*

GUARANTEED ANALYSIS

Crude protein equivalent from NPN¹ (max)..... %
Crude Fat, not less than.....%
Crude Fiber, not more than.....%
Calcium (min)..... %
Calcium (max)..... %
Phosphorus (min)..... %
Salt¹ (min)..... %
Salt¹ (max)..... %
Sodium² (min)..... %
Sodium² (max)..... %
Potassium (min)..... %
Vitamin A^{1,3}..... IU/LB

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

³Other than precursors of Vitamin A, if added

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Blend with grains and non-medicated feed ingredients to provide a complete feed with the following grams of chlortetracycline per ton in the resulting Type C medicated feed.

Indications for Use

**Use Levels of
Chlortetracycline**

CATTLE:

For Growing Cattle (over 400 lbs): Mix an amount of Type B feed to provide approximately 70 mg/head/day varying with body weight and feed consumption with non-medicated feed ingredients in the resulting Type C medicated feed to make the daily complete feed for the reduction of the incidence of liver abscesses.

For Beef Cattle: Mix an amount of Type B feed to provide approximately 350 mg/head/day varying with body weight and feed consumption with non-medicated feed ingredients in the resulting Type C medicated feed to make the daily complete feed for the control of bacterial pneumonia associated with shipping fever complex *Pasteurella* spp. susceptible to chlortetracycline.

For Beef Cattle (under 700 lbs): Mix an amount of Type B feed to provide approximately 350 mg/head/day varying with body weight and feed consumption with non-medicated feed ingredients in the resulting Type C medicated feed to make the daily complete feed for the control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline.

350 mg/head/day

For Beef Cattle (over 700 lbs): Mix an amount of Type B feed to provide approximately 0.5 mg/lb body weight/day varying with body weight and feed consumption with non-medicated feed ingredients in the resulting Type C medicated feed to make the daily complete feed for the control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline.

0.5 mg/lb body weight/day

For Calves, Beef and Nonlactating Dairy Cattle: Mix an amount of Type B feed to provide approximately 10 mg/lb body weight/day varying with body weight and feed consumption with non-medicated feed ingredients in the resulting Type C medicated feed to make the daily complete feed for the treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to chlortetracycline.
(Treat for not more than 5 days)

10 mg/lb body weight/day

WARNING: A WITHDRAWAL PERIOD HAS NOT BEEN ESTABLISHED FOR THIS PRODUCT IN PRE-RUMINATING CALVES. DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL. ZERO-DAY WITHDRAWAL PERIOD.

*The final printed label must bear a single drug concentration.

Net Weight _____ lb (_____ kg)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000