

**Chlortetracycline Chicken Supplement
Type B Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated chicken feed.

ACTIVE DRUG INGREDIENT

Chlortetracycline.....2-40 g/lb*

GUARANTEED ANALYSIS

Crude Protein, not less than.....%
Crude Fat, not less than.....%
Crude Fiber, not more than.....%
Lysine (min).....%
Methionine (min).....%
Calcium (min).....%
Calcium (max).....%
Phosphorus(min).....%
Salt¹ (min).....%
Salt¹ (max).....%
Sodium² (min).....%
Sodium² (max).....%

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Blend with grains and non-medicated feed ingredients to provide a complete feed with the following grams of chlortetracycline per ton of complete feed.

Indications for Use

**Use Levels of
Chlortetracycline**

CHICKENS

For Chickens: Mix 50 to 100 pounds of Type B (2 g/lb) or 2.5 to 5 pounds of Type B (40 g/lb) with non-medicated feed ingredients to make a ton of broiler chicken feed for the control of infectious synovitis caused by *Mycoplasma synoviae*, susceptible to chlortetracycline containing 100 to 200 g/ton to be fed continuously for 7 to 14 days.

WARNING: Do not feed to chickens producing eggs for human consumption. Zero-day withdrawal period.

100-200 g/ton

For Chickens: Mix 100 to 200 pounds of Type B (2 g/lb) or 5 to 10 pounds of Type B (40 g/lb) with non-medicated feed ingredients to make a ton of chickens feed for the control of chronic respiratory disease (CRD) and air sac infection caused by *Mycoplasma gallisepticum* and *Escherichia coli* susceptible to chlortetracycline containing 200 to 400 g/ton to be fed

200-400 g/ton

continuously for 7 to 14 days.

WARNING: Do not feed to chickens producing eggs for human consumption. Zero-day withdrawal period.

For Chickens: Mix 250 pounds of Type B (2 g/lb) or 12.5 pounds of Type B (40 g/lb) with non-medicated feed ingredients to make a ton of chickens feed for the reduction of mortality due to *Escherichia coli* infection susceptible to chlortetracycline containing 500 g/ton to be fed continuously for 5 days. 500 g/ton

WARNING: Do not feed to chickens producing eggs for human consumption. Withdraw 24 hours prior to slaughter.

*The final printed label must bear a single drug concentration.

Net Weight _____ lb (_____ kg)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000