

**Oxytetracycline Chicken Supplement
Type B Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated chicken feed.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to *2-20 grams oxytetracycline hydrochloride/lb.

GUARANTEED ANALYSIS

Crude Protein, (Min).....	XX.XX %
Lysine, (Min).....	X.XX%
Methionine, (Min).....	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min).....	X.XX%
Calcium, (Max).....	X.XX%
Phosphorus, (Min).....	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%
Potassium, (Min) ¹	X.XX%

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below. To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing as a complete feed.

Indications for Use

Oxytetracycline in
Type C medicated feed

CHICKENS

For Chickens: Mix 50 to 100 pounds of Type B (2 g/lb) or 5 to 10 pounds of Type B (20 g/lb) with non-medicated feed ingredients to make a ton of chicken feed for control of infectious synovitis caused

100-200 g/ton

by Mycoplasma synoviae and control of fowl cholera caused by Pasteurella multocida susceptible to oxytetracycline containing 100 to 200 g/ton to be fed continuously for 7 to 14 days.

RESIDUE WARNING: Do not feed to chickens producing eggs for human consumption. Zero-day withdrawal period. Do not use in low-calcium feed containing less than 0.55% dietary calcium. Use in such feeds may result in violative residues.

For Chickens: Mix 200 pounds of Type B (2 g/lb) or 20 pounds of Type B (20 g/lb) with non-medicated feed ingredients to make a ton of chicken feed for control of chronic respiratory disease (CRD) and air sac infection caused by Mycoplasma gallisepticum and Escherichia coli susceptible to oxytetracycline containing 400 g/ton to be fed continuously for 7 to 14 days. 400 g/ton

RESIDUE WARNING: Do not feed to chickens producing eggs for human consumption. Zero-day withdrawal period. Do not use in low-calcium feed containing less than 0.55% dietary calcium. Use in such feeds may result in violative residues.

CHICKENS

For Broiler Chickens: Mix 250 pounds of Type B (2 g/lb) or 25 pounds of type B (20 g/lb) with non-medicated feed ingredients to make a ton of chicken feed for reduction of mortality due to air sacculitis (air-sac-infection) caused by Escherichia coli susceptible to oxytetracycline containing 500 g/ton to be fed continuously for 5 days. 500 g/ton

RESIDUE WARNING: Do not feed to chickens producing eggs for human consumption. 24 hour withdrawal period. Do not use in low-calcium feed containing less than 0.55% dietary calcium. Use in such feeds may result in violative residues.

*The medicated feed label must state a single drug concentration.

Net Weight _____ lb (_____ kg)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000