BLUE BIRD LABELING

chlortetracycline hydrochloride

Type C Medicated Turkey Feed – Bluecomb

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline in turkeys.

Active Drug Ingredient	
Chlortetracycline* mg/lb (e	;/ton)**
Guaranteed Analysis	
Crude Protein (Min)	%
Lysine (Min)	
Methionine (Min)	%
Crude Fat (Min)	%
Crude Fiber (Max)	
Calcium (Min)	%
Calcium (Max)	%
Phosphorus (Min)	%
Salt ^a (Min)	%
Salt ^a (Max)	
Sodium ^b (Min)	
Sodium ^b (Max)	%
^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal soul b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.	ce of the nutrient.
Ingredients	
Ingredients as defined by AAFCO.	
Feeding Directions	
Feed continuously for 7 to 14 days, providing chlortetracycline at the rate of 25 mg per pound of b day. ()*** pounds of this feed will medicate 100 lbs of body weight.	oody weight per
Warning	

Do not feed to turkeys producing eggs for human consumption.

Manufactured By:

Blue Bird Feed Mill City, State, Zip

NET WEIGHT ON BAG OR BULK

- *Deracin is the trade name of chlortetracycline hydrochloride formulation (Type A Medicated Article).
- **Label must show actual concentration of chlortetracycline, which will vary with body weight and feed consumption to provide 25 mg/lb body weight per day.
- *** The following table provides examples of feeding rates and amount of drug per ton of feed. Third column represents number of pounds of feed that will medicate 100 lbs of body weight.

Drug level in feed		** Ibs of Turkey Feed
mg/lb	grams/ton	Per 100 lbs Body Weight
500	1000	5.0
625	1250	4.0
833	1667	3.0
1250	2500	2.0
2500	5000	1.0