# **Product Profile**

# CircoGard®

## Porcine Circovirus Vaccine, Type 2, Killed Baculovirus Vector

### **Product Description**

- One-dose porcine circovirus (PCV) Type 2b vaccine for vaccination of swine
- Inactivated trivalent-adjuvanted formulation using StimGard Plus™ adjuvant
- Each vial contains 50 or 250 doses (1 mL/dose)



#### **Advantages**

- Labor-saving 1-mL dose easy on nursing piglets and staff.
- Safe for use in young animals.
- Demonstrated heterologous circovirus protection.
- 3 different classes of adjuvants in StimGard Plus<sup>™</sup> help provide for an optimal immune response.

#### Dosage / Administration

 Shake well before use. Using aseptic technique, inject a single 1.0-mL dose intramuscularly (IM).

#### **Indications / Uses**

- For use in healthy piglets of all breeds, 10 days of age or older, against Porcine Circovirus, Type
   2.
- Visit USDA website for full label indications & technical information. (productdata.aphis.usda.gov).

#### **Precautions**

- CAUTION: For animal use only.
- Store at 2 to 8°C (35-45°F). Do not freeze. Shake well before use. Use entire contents upon opening.
- Do not vaccinate within 21 days before slaughter.
- Transient reaction may occur at the injection site. If anaphylaxis occurs, use epinephrine or equivalent. Contains merthiolate (Thimerosal) as a preservative.
- This product has not been tested in pregnant animals.

#### **Packaging**

- 50-mL or 250-mL vials.
- Other sizes available.

#### **Ordering Information**

• Prod. number: 22010-50 (50 mL) 22010-250 (250 mL)

## **Key Features**

- Pig-proven under field conditions.
- StimGard Plus<sup>™</sup> trivalent adjuvant system helps optimize both cellmediated and humoral immune responses.
- Safe.
- Excellent syringeability.
- Advanced purification process allows low-volume injection for minimal tissue reaction.
- QR (quick response) codes on the label for inventory and tracking purposes.



# CircoGard

## Porcine Circovirus Vaccine, Type 2, Killed Baculovirus Vector

Porcine Circovirus Vaccine, Type 2, Killed Baculovirus Vector

CircoGard®



This product has been shown to be effective for the vaccination of healthy piglets, 10 days of age or older against Porcine Circovirus, Type 2. To review efficacy and safety data, go to productdata.aphis.usda.gov.

ADMINISTRATION AND DOSAGE:

Dose: 1.0 mL intramuscularly (I.M.).

CAUTION: For animal use only. Do not mix with other products, except as specified on this label. In case of human exposure, contact a physician. Store at 2-8°C (35-45°F). Do not freeze. Shake well before use. Use entire contents upon opening. Do not vaccinate within 21 days before slaughter. Transient reaction may occur at injection site. If anaphylaxis occurs, use epinephrine or equivalent. Contains merthiolate (Thimerosal) as a preservative. This product has not been tested in pregnant animals. Revaccination interval not established.

**Product No:** Exp. Date:

Manufactured by:

**Pharmgate Biologics** 2575 University Avenue West, Suite 100, St. Paul. MN 55114 612-256-0930

VLN 329 / PCN 19K5.R0

Visit USDA website for full label indications & technical information: productdata.aphis.usda.gov

Pharmgate Animal Health swine vaccines/bacterins.		
Product	Doses	Pharmgate Product Number
MycoGard® (2 dose)	50	20020-50
MycoGard® (2 dose)	250	20020-250
MycoGard-1™ (1 dose)	50	21010-50
MycoGard-1™ (1 dose)	250	21010-250
CircoGard® (1 dose)	50	22010-50
CircoGard® (1 dose)	250	22010-250
Circo/MycoGard® (1 dose)	50	25010-50
Circo/MycoGard® (1 dose)	250	25010-250

