

CTC / MONENSIN 10mg/BE-BP FE – BEEF FEEDLOT CATTLE FEED
(chlortetracycline and monensin Type C medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Indications for Use

For treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to chlortetracycline and for improved feed efficiency in growing beef steers and heifers fed in confinement for slaughter.

Active Drug Ingredients

Chlortetracycline^a (as chlortetracycline calcium complex) equivalent to chlortetracycline hydrochloride..... 400 to 2000 g/ton*
Monensin, USP^b..... 5 to 40 g/ton*

Guaranteed Analysis

Crude protein (min)	_____	%
NPN ¹ (max)	_____	%
Crude fat (min)	_____	%
Crude fiber (max)	_____	%
Calcium ¹ (min)	_____	%
Calcium ¹ (max)	_____	%
Phosphorus ¹ (min)	_____	%
Salt ¹ (min)	_____	%
Salt ¹ (max)	_____	%
Sodium ² (min)	_____	%
Sodium ² (max)	_____	%
Potassium ¹ (min)	_____	%
Vitamin A ¹ (min)	_____	IU/lb

¹ Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of nutrient.

²Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient as named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

Feeding Directions

Feed as the sole ration to provide 10 mg chlortetracycline per lb. body weight per day and 50 to 480 mg monensin per head per day. Feed for not more than 5 days, then continue feeding monensin Type C medicated feed alone. No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 grams per ton (360 mg monensin per head per day).

The following table provides examples of feeding rates and amount of drug per ton of feed:

Body Weight (lbs)	Feed Consumption		Chlortetracycline in medicated feed		Desired Intake of Monensin	Monensin in medicated feed	
	% body weight	lb/head/day	mg/lb	g/ton	mg/hd/day	mg/lb	g/ton
1000	1.5	15	666.67	1,333.33	150	10	20
	2	20	500	1,000	400	20	40
	3	30	333.33	666.67	75	2.5	5
1200	1.5	18	666.67	1,333.33	180	10	20
	2	24	500	1,000	480	20	40
	3	36	333.33	666.67	90	2.5	5
1400	1.5	21	666.67	1,333.33	420	20	40
	2	28	500	1,000	280	10	20
	3	42	333.33	666.67	105	2.5	5

Caution

For use in dry feeds only. Not for use in liquid feed supplements. Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning

Withdrawal Periods and Residue Warnings:

No withdrawal period is required when used according to labeling. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

User Safety Warnings:

Keep this and all drugs out of the reach of children. Not for human use.

Lot No. (if applicable) _____

Approved by FDA under NADA # 141-564

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000

NET WEIGHT ON BAG OR BULK

*Final printed label on formulated Type C medicated feed must bear a single drug concentration.

^a Sourced from Pennchlor[®], NADA # 138-935

^b Sourced from Rumensin[™], NADA # 95-735