

**Bacitracin Methylenedisalicylate and Monensin Broiler Chicken Feed – ROWG/FE**  
 (bacitracin methylenedisalicylate and monensin Type C medicated feed)

**INDICATIONS FOR USE**

For increased rate of weight gain and improved feed efficiency, and as an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima* in broiler chickens.

**ACTIVE DRUG INGREDIENTS**

Bacitracin (as feed grade bacitracin methylenedisalicylate)<sup>a</sup>..... 4 to 50 g/ton\*  
 Monensin,<sup>b</sup> USP..... 90 to 110 g/ton\*

**GUARANTEED ANALYSIS**

Crude protein, not less than	.....	_____	%
Lysine, not less than	.....	_____	%
Methionine, not less than	.....	_____	%
Crude fat, not less than	.....	_____	%
Crude fiber, not more than	.....	_____	%
Calcium, not less than	.....	_____	%
Calcium, not more than	.....	_____	%
Phosphorus, not less than	.....	_____	%
Salt <sup>1</sup> , not less than	.....	_____	%
Salt <sup>1</sup> , not more than	.....	_____	%
Sodium <sup>2</sup> , not less than	.....	_____	%
Sodium <sup>2</sup> , not more than	.....	_____	%

<sup>1</sup> If added

<sup>2</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**FEEDING DIRECTIONS**

Feed as the sole ration throughout the feeding period.

**CAUTION**

Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age. Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. In the absence of coccidiosis in broiler chickens the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain. Not for broiler breeder replacement chickens.

**WARNING**



**Withdrawal Periods:** No withdrawal period is required when used according to labeling.



**Questions/Comments?** Contact Pharmgate Animal Health LLC at 1-800-380-6099 or [www.pharmgate.com](http://www.pharmgate.com). To report side effects, contact Pharmgate Animal Health LLC at 1-833-531-0114. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

Approved by FDA under NADA # 141-565

**MANUFACTURED BY**

Blue Bird Feed Company  
Robin, Indiana 11111

**NET WEIGHT ON BAG OR BULK**

Lot Number (if applicable): \_\_\_\_\_

Do not use this medicated feed after: \_\_\_\_\_  
(90 days from the date of manufacture)

\*The final printed feed label should list only a single drug concentration.

<sup>a</sup>Pennitracin MD 50G<sup>®</sup> is the proprietary name of the bacitracin methylenedisalicylate formulation (Type A medicated article) approved by FDA under NADA # 141-137.

<sup>b</sup>Coban<sup>™</sup> is the proprietary name of the monensin formulation (Type A medicated article) approved by FDA under NADA # 38-878.