

**Bacitracin Methylenedisalicylate and Narasin and Nicarbazin Complete Feed
for Broiler Chickens – NE-P**
(bacitracin methylenedisalicylate and narasin and nicarbazin Type C medicated feed)

INDICATIONS FOR USE

For the prevention of mortality caused by necrotic enteritis associated with *Clostridium perfringens*, and for the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima* in broiler chickens.

ACTIVE DRUG INGREDIENTS

Bacitracin (as feed grade bacitracin methylenedisalicylate) ^a	50 g/ton
Narasin ^b	27 to 45 g/ton*
Nicarbazin ^b	27 to 45 g/ton*

GUARANTEED ANALYSIS

Crude protein, not less than	_____	%
Lysine, not less than	_____	%
Methionine, not less than	_____	%
Crude fat, not less than	_____	%
Crude fiber, not more than	_____	%
Calcium, not less than	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than	_____	%
Salt ¹ , not less than	_____	%
Salt ¹ , not more than	_____	%
Sodium ² , not less than	_____	%
Sodium ² , not more than	_____	%

¹ If added

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed as the sole ration for 28 to 35 days, starting from the time chicks are placed for brooding.

CAUTION

For broiler chickens only. Do not feed to laying hens. Do not allow adult turkeys, horses or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal. Nicarbazin medicated broilers may show reduced heat tolerance if exposed to high temperature and high humidity. Provide adequate drinking water and ventilation during these periods.

WARNING



Withdrawal Periods: No withdrawal period is required when used according to this labeling.



Questions/Comments? Contact Pharmgate Animal Health LLC at 1-800-380-6099 or www.pharmgate.com. To report side effects, contact Pharmgate Animal Health LLC at 1-833-531-0114. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

Approved by FDA under NADA # 141-529

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

Lot Number (if applicable): _____

*The final printed feed label should list only a single drug concentration. A fixed ratio (1:1) Type A medicated article of narasin and nicarbazin (Maxiban™) has to be used in the manufacture of this combination.

^aPennitracin MD 50G® is the proprietary name of the bacitracin methylenedisalicylate formulation (Type A medicated article) approved by FDA under NADA # 141-137.

^bMaxiban™ is the proprietary name of the narasin and nicarbazin formulation (Type A medicated article) approved by FDA under NADA # 138-952.