

Product Profile


Pennox 50[®]



Oxytetracycline Type A Medicated Article

Product Description

- Antibacterial premix for oral administration in feed to chickens, turkeys, cattle, swine, sheep, and bees
- Each pound of premix contains 50 grams of oxytetracycline activity



Formulation

- Free-flowing meal feed medication.
- Ingredients: Oxytetracycline dihydrate base, calcium carbonate, roughage products, mineral oil.

FDA Status

- CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.
- Type A Medicated Article for use in the manufacture of medicated dry feeds (not for use in liquid feeds).
- Category I drug; does not require a feedmill license.

Withdrawal Period

- 0 days for swine and turkeys.
- 0-5 days for chickens and cattle.
- Different withdrawal times may be required for certain export markets.

Indications (Partial List, See Back)

- Chickens: Control of infectious synovitis (*Mycoplasma synoviae*); control of fowl cholera (*Pasteurella multocida*); control of chronic

respiratory disease (CRD) and air sac infection (*Mycoplasma gallisepticum*, *Escherichia coli*); reduction of mortality due to air sacculitis (*E. coli*).

- Turkeys: Control of infectious synovitis (*M. synoviae*); control of hexamitiasis (*Hexamita meleagrides*); control of complicating bacteria associated with bluecomb (transmissible enteritis, coronaviral enteritis).
- Cattle: Reduction of incidence of liver abscesses; prevention and treatment of bacterial pneumonia/shipping fever complex (*P. multocida*); treatment of bacterial enteritis (*E. coli*).
- Swine: Treatment of bacterial enteritis (*E. coli*, *Salmonella choleraesuis*) and bacterial pneumonia (*P. multocida*); control and treatment of leptospirosis (*Leptospira pomona*); reducing incidence of abortion and shedding of leptospirae).

Packaging

- 50-lb, red color-coded bags.

Key Features

- Convenient, cost-effective treatment option vs injectable therapy.
- Tried and trusted.
- Broad spectrum, effective against both Gram-positive and Gram-negative organisms that can cause respiratory, enteric, or reproductive diseases.
- Readily absorbed, providing effective blood and lung tissue concentrations.
- Free-flowing formulation promotes consistent mixing.
- Wide safety margin.
- Versatile, with wide range of approved combinations.
- Color-coded bag for easy identification.
- High-quality manufacturing at US Pharmgate facilities according to FDA requirements.

The labeling contains complete use information, including any cautions and warnings. Always read, understand and follow the labeling and use directions. See the reverse side for use directions and additional information.



Pharmgate
ANIMAL HEALTH

1800 Sir Tyler Drive
Wilmington, NC | 28405
910.679.8364
Pharmgate.com

Pennox 50[®]

Oxytetracycline
Type A Medicated Article

Active Drug Ingredients:		Oxytetracycline (from oxytetracycline dihydrate base) equivalent to 50 grams Oxytetracycline Hydrochloride/lb.		
Ingredients:		Oxytetracycline, Calcium Carbonate, Roughage Products and Mineral Oil.		
For Use in the Manufacture of Medicated Feeds. Do Not Feed Undiluted.				
CAUTION: For use in Dry Feeds ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS				
Directions For Use				
Indications For Use	Use Levels of Oxytetracycline	lbs. of Pennox 50[®] per ton	Indications For Use	
Use Levels of Oxytetracycline		lbs. of Pennox 50[®] per ton		
<p>Chickens</p> <p>For Chickens: Control of infectious synovitis caused by <i>Mycoplasma synoviae</i>; control of fowl cholera caused by <i>Pasteurella multocida</i> sensitive to oxytetracycline. (Feed continuously for 7-14 days)</p> <p>For Chickens: Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to oxytetracycline. (Feed continuously for 7-14 days)</p> <p>For Broiler Chickens: Reduction of mortality due to air sacculitis (air sac infection) caused by <i>Escherichia coli</i> susceptible to oxytetracycline. (Feed continuously for 5 days)</p> <p>Withdrawal Periods and Residue Warnings: Do not feed to chickens producing eggs for human consumption. Do not use in low calcium feed containing less than 0.55% dietary calcium. Use in such feeds may result in violative residues. 24 hour withdrawal period at 500 g/ton use level. No withdrawal period is required when used according to labeling at 100-200 g/ton and 400 g/ton use levels.</p>	<p>100-200g/ton</p> <p>400g/ton</p> <p>500g/ton</p>	<p>2.0-4.0</p> <p>8.0</p> <p>10.0</p>	<p>Calves, Beef Cattle, and Nonlactating Dairy Cattle</p> <p>For Growing Cattle (over 400 lbs.): For the reduction of the incidence of liver abscesses. (Use continuously)</p> <p>For Cattle: For the prevention and treatment of the early stages of the shipping fever complex. (Feed 3-5 days before and after arrival in feedlots.)</p> <p>For Calves, Beef Cattle and Nonlactating Dairy Cattle: Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline. (Feed continuously for 7-14 days)</p> <p>Withdrawal Periods and Residue Warnings: This drug product is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. 5-day withdrawal period at 10 mg/lb. body weight/day use level. No withdrawal period is required when used according to labeling at 75 mg/head/day and 0.5-2.0 g/head/day use levels.</p>	<p>75 mg/head/day</p> <p>0.5-2.0g/head/day</p> <p>10 mg/lb. body weight/day</p>
<p>Turkeys</p> <p>For Turkeys: Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to oxytetracycline. (Feed continuously for 7-14 days)</p> <p>For Turkeys: Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to oxytetracycline. (Feed continuously for 7-14 days)</p> <p>For Turkeys: Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline. (Feed continuously for 7-14 days)</p> <p>Withdrawal Periods and Residue Warnings: Do not feed to turkeys producing eggs for human consumption. Zero-day withdrawal period.</p>	<p>100g/ton</p> <p>200g/ton</p> <p>25 mg/lb. body weight/day</p>	<p>2.0</p> <p>4.0</p>	<p>Honey Bees</p> <p>Control of American Foulbrood caused by <i>Paenibacillus larvae</i>, and European Foulbrood caused by <i>Melissococcus plutonius</i> susceptible to oxytetracycline.</p> <p>200 mg/oz (Dusting) 200 mg/5 lb (Syrup) 800 mg/patty (Extender Patty)</p> <p>See Mixing, Preparation, and Feeding Directions below.</p> <p>Mixing Directions for Honey Bees</p> <p>To manufacture a Type B medicated feed containing 20 g/lb of oxytetracycline hydrochloride activity, mix 10 lb of Pennox 50 with 15 lb of powdered sugar (for a total of 25 lb.)</p> <p>To manufacture a Type C medicated feed containing 200 mg of oxytetracycline hydrochloride activity per oz (dusting Type C medicated feed), mix 14 lb of Pennox 50 with 204 lb of powdered sugar (for a total of 218 lb). The resulting Type C medicated feed is fed as-is for dusting, or is used to prepare syrup and extender patty feeds.</p> <p>Feeding Directions for Dusting:</p> <p>Dusting: Apply 1 oz (200 mg oxytetracycline) of dusting Type C medicated feed per colony every 4 to 5 days for a total of 3 applications. Apply the dust on the outer parts or ends of the frames.</p> <p>Caution: Dusting of uncapped brood cells has been reported to cause death of larval honey bees. Do not dust uncapped brood cells.</p> <p>Preparation and Feeding Directions for Syrup:</p> <p>Preparation: First dissolve 1 oz (200 mg oxytetracycline) of the dusting Type C medicated feed in a small quantity of water, then mix into 5 lb of 1:1 sugar syrup (equal parts sugar and water, w/w).</p> <p>Feeding: Feed 5 lb of syrup (200 mg oxytetracycline) per colony every 4 to 5 days for a total of 3 applications. Bulk feed the syrup using feeder pails, or division board feeders, or by filling the combs. Prepare fresh syrup for each application.</p> <p>Preparation and Feeding Directions for Extender Patty:</p> <p>Preparation: Make one patty by mixing 4 oz (800 mg oxytetracycline) of the dusting Type C medicated feed with 5.8 oz (165 g) of vegetable shortening and 11.6 oz (330 g) of sugar.</p> <p>Feeding: Place one patty per colony on the top bars of the brood nest frames in a single application.</p> <p>Warnings:</p> <p>Withdrawal Periods and Residue Warnings: Remove at least 6 weeks prior to main honey flow. Type C medicated feeds should be fed in the spring or fall and consumed by the bees before main honey flow begins to avoid contamination of production honey. Honey stored during medication periods in combs for surplus honey should be removed following final medication of the bee colony and must not be used for human food.</p> <p>Warning: Do not use in a manner contrary to state apiaary laws and regulations. Each state has specific regulations relative to disease control and medication. Contact the appropriate official or state departments of agriculture for specific inter- and intrastate laws and regulations.</p>	
<p>Swine</p> <p>For Swine: Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> susceptible to oxytetracycline and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline. (Feed continuously for 7-14 days)</p> <p>For Breeding Swine: Control and treatment of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to oxytetracycline. (Feed continuously for 14 days)</p> <p>Withdrawal Periods: Zero-day withdrawal period.</p>	<p>Feed approximately 400g/t, varying with body weight and feed consumption to provide 10mg/lb. body weight/day</p> <p>Feed approximately 400g/t, varying with body weight and feed consumption to provide 10mg/lb. body weight/day</p>			
<p>Sheep</p> <p>For Sheep: Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline. (Feed continuously for 7-14 days)</p> <p>Withdrawal Periods: 5-day withdrawal period.</p>	<p>10 mg/lb. body weight/day</p>			
<p>Approved by FDA under NADA # 138-938 Pennox 50[®] is the registered trademark of Pharmgate Inc.</p>				

