

**CHICKENS**  
**(oxytetracycline Type B medicated feed)**

**Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

For use in manufacturing medicated chicken feed.

**Indications for Use:** Control of infectious synovitis caused by *Mycoplasma synoviae* and control of fowl cholera caused by *Pasteurella multocida* sensitive to oxytetracycline in chickens, control of chronic respiratory disease (CRD) and air sac infection caused by *Mycoplasma gallisepticum* and *Escherichia coli* susceptible to oxytetracycline in chickens, and reduction of mortality due to air sacculitis (air sac infection) caused by *Escherichia coli* susceptible to oxytetracycline in broiler chickens.

**ACTIVE DRUG INGREDIENT**

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to oxytetracycline hydrochloride<sup>a</sup>.....2-20 g/lb\*

**GUARANTEED ANALYSIS**

Crude Protein (Min) .....	_____	%
Lysine (Min).....	_____	%
Methionine (Min) .....	_____	%
Crude Fat (Min).....	_____	%
Crude Fiber (Max).....	_____	%
Calcium (Min).....	_____	%
Calcium (Max) .....	_____	%
Phosphorus (Min).....	_____	%
Salt <sup>1</sup> (Min).....	_____	%
Salt <sup>1</sup> (Max) .....	_____	%
Sodium <sup>2</sup> (Min).....	_____	%
Sodium <sup>2</sup> (Max) .....	_____	%

<sup>1</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>2</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Ingredients as defined by AAFCO

**MIXING DIRECTIONS**

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below. To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing as a complete feed.

Indications for Use

Oxytetracycline in  
Type C medicated feed

For Chickens: Mix 50 to 100 pounds of Type B (2 g/lb) or 5 to 10 pounds of Type B (20 g/lb) with non-medicated feed ingredients to make a ton of chicken feed for control of infectious synovitis caused by *Mycoplasma synoviae* and control of fowl cholera caused by *Pasteurella multocida* susceptible to oxytetracycline containing 100 to 200 g/ton to be fed continuously for 7 to 14 days.

100-200 g/ton

For Chickens: Mix 200 pounds of Type B (2 g/lb) or 20 pounds of Type B (20 g/lb) with non-medicated feed ingredients to make a ton of chicken feed for control of chronic respiratory disease (CRD) and air sac infection caused by *Mycoplasma gallisepticum* and *Escherichia coli* susceptible to oxytetracycline containing 400 g/ton to be fed continuously for 7 to 14 days.

400 g/ton

For Broiler Chickens: Mix 250 pounds of Type B (2 g/lb) or 25 pounds of Type B (20 g/lb) with non-medicated feed ingredients to make a ton of chicken feed for reduction of mortality due to air sacculitis (air-sac-infection) caused by *Escherichia coli* susceptible to oxytetracycline containing 500 g/ton to be fed continuously for 5 days.

**Withdrawal Periods and Residue Warnings:** Do not feed to chickens producing eggs for human consumption. Do not use in low calcium feed containing less than 0.55% dietary calcium. Use in such feeds may result in violative residues. 24 hour withdrawal period at 500 g/ton use level. No withdrawal period is required when used according to labeling at 100-200 g/ton and 400 g/ton use levels.

Net Weight \_\_\_\_\_ lb (\_\_\_\_\_ kg)

Approved by FDA under NADA # 138-938  
Manufactured by  
Blue Bird Feed Company  
Blue Bird, MD 00000

Lot number (if applicable) \_\_\_\_\_

\*The final printed medicated feed label must state a single drug concentration.

<sup>a</sup> Pennox<sup>®</sup> is the proprietary name of oxytetracycline Type A medicated article (NADA # 138-938).

Pharmgate Inc. [June 24, 2022]