

**SWINE
(oxytetracycline Type B medicated feed)**

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated swine feed.

Indications for Use: Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* susceptible to oxytetracycline and treatment of bacterial pneumonia caused by *Pasteurella multocida* susceptible to oxytetracycline in swine and control and treatment of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by *Leptospira pomona* susceptible to oxytetracycline in breeding swine.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to oxytetracycline hydrochloride^a.....2-20 g/lb*

GUARANTEED ANALYSIS

Crude Protein (Min)	_____ %
Lysine (Min).....	_____ %
Crude Fat (Min).....	_____ %
Crude Fiber (Max).....	_____ %
Calcium (Min).....	_____ %
Calcium (Max)	_____ %
Phosphorus (Min).....	_____ %
Salt ¹ (Min).....	_____ %
Salt ¹ (Max)	_____ %
Sodium ² (Min).....	_____ %
Sodium ² (Max)	_____ %
Selenium ³ (Min)	_____ ppm

¹ Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

² Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

³ Guarantee shall be stated in ppm when less than 10,000 ppm and in percentage when concentration is 10,000 ppm (1%) or greater.

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below. To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing as a complete feed.

Indications for Use

For Swine: Mix an amount of Type B feed to provide approximately 400 g/ton varying with body weight and feed consumption to provide 10 mg/lb body weight/day with non-medicated feed ingredients to make a ton of swine feed for treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to oxytetracycline to be fed continuously for 7 to 14 days.

For Breeding Swine: Mix an amount of Type B feed to provide approximately 400 g/ton varying with body weight and feed consumption to provide 10 mg/lb body weight/day with non-medicated feed ingredients to make a ton of swine feed for



**Oxytetracycline in
Type C medicated feed**

Feed approximately 400 g/ton, varying with body weight and feed consumption, to provide 10 mg/lb body weight/day

Feed approximately 400 g/ton, varying with body weight and feed consumption, to provide

control and treatment of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by *Leptospira pomona* susceptible to oxytetracycline to be fed continuously for 14 days.

10 mg/lb body weight/day

 **Withdrawal Periods:** Zero-day withdrawal period. 

Net Weight _____ lb (_____ kg)

Approved by FDA under NADA # 138-938
Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000

Lot number (if applicable) _____

*The final printed medicated feed label must state a single drug concentration.

^a Pennox[®] is the proprietary name of oxytetracycline Type A medicated article (NADA # 138-938).

Pharmgate Inc. [June 24, 2022]