

TURKEYS

(oxytetracycline Type B medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated turkey feed.

Indications for Use: Control of hexamitiasis caused by *Hexamita meleagridis* susceptible to oxytetracycline, control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to oxytetracycline, and control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline in turkeys.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to oxytetracycline hydrochloride^a.....2-20 g/lb*

GUARANTEED ANALYSIS

Crude Protein (Min)	_____	%
Lysine (Min).....	_____	%
Methionine (Min)	_____	%
Crude Fat (Min).....	_____	%
Crude Fiber (Max).....	_____	%
Calcium (Min).....	_____	%
Calcium (Max)	_____	%
Phosphorus (Min).....	_____	%
Salt ¹ (Min).....	_____	%
Salt ¹ (Max)	_____	%
Sodium ² (Min).....	_____	%
Sodium ² (Max)	_____	%

¹ Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

² Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below. To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing as a complete feed.

Indications for Use	Oxytetracycline in Type C medicated feed
For Turkeys: Mix 50 pounds of Type B (2 g/lb) or 5 pounds of Type B (20 g/lb) with non-medicated feed ingredients to make a ton of turkey feed for control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to oxytetracycline containing 100 g/ton to be fed continuously for 7 to 14 days.	100 g/ton
For Turkeys: Mix 100 pounds of Type B (2 g/lb) or 10 pounds of Type B (20 g/lb) with non-medicated feed ingredients to make a ton of turkey feed for control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to oxytetracycline containing 200 g/ton to be fed continuously for 7 to 14 days.	200 g/ton
For Turkeys: Mix an amount of Type B feed to provide approximately 25 mg/lb body weight/day varying with body weight and feed consumption with non-medicated ingredients to make a ton of turkey feed for control of complicating bacterial organisms associated with bluecomb (transmissible	25 mg/lb body weight/day

enteritis, coronaviral enteritis) susceptible to oxytetracycline to be fed continuously for 7 to 14 days.



Withdrawal Periods and Residue Warnings: Do not feed to turkeys producing eggs for human consumption. Zero-day withdrawal period.



Net Weight _____ lb (_____ kg)

Approved by FDA under NADA # 138-938
Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000

Lot number (if applicable) _____

*The final printed medicated feed label must state a single drug concentration.

^a Pennox[®] is the proprietary name of oxytetracycline Type A medicated article (NADA # 138-938).

Pharmgate Inc. [June 24, 2022]