



MycoGard-1™

Mycoplasma hyopneumoniae Bacterin

Product Description

- One-dose *Mycoplasma hyopneumoniae* bacterin for vaccination of swine
- Inactivated trivalent-adjuvanted formulation using StimGard Plus™ adjuvant
- Each vial contains 50 or 250 doses (1 mL/dose)



Advantages

- 1-mL dose – easy dose size for nursing piglets.
- Convenient dosing – allowing for very early *Mycoplasma hyopneumoniae* vaccination.
- Safe – easy on animals.
- 3 different classes of adjuvants in StimGard Plus™ help provide for an optimal immune response.
- Colored cap - for ease of identification.

Dosage / Administration

- Shake well before use. Using aseptic technique, inject a single 1.0-mL dose intramuscularly (IM).

Indications / Uses

- For use in healthy piglets of all breeds, 10 days of age or older, against *Mycoplasma hyopneumoniae*.
- Visit USDA website for full label indications & technical information. (productdata.aphis.usda.gov).

Precautions

- CAUTION: For animal use only.
- Store at 2 to 8°C (35-45°F). Do not freeze. Shake well before use. Use entire contents upon opening.
- Do not vaccinate within 21 days before slaughter.
- Transient reaction may occur at the injection site. If anaphylaxis occurs, use epinephrine or equivalent. Contains merthiolate (Thimerosal) as a preservative.
- This product has not been tested in pregnant animals.

Packaging

- 50-mL or 250-mL vials.

Ordering Information

- Prod. number:
21010-50 (50 mL)
21010-250 (250 mL)

Key Features

- StimGard Plus™ trivalent adjuvant system helps optimize both cell-mediated and humoral immune responses.
- Excellent syringeability.
- Advanced purification process allows low-volume injection for minimal tissue reaction.
- Cap colored for easy identification.
- QR (quick response) codes on the label for inventory and tracking purposes.

The labeling contains complete use information, including cautions and warnings. Always read, understand and follow the labeling and use directions. See the back page for use directions and additional information.

Mycogard-1™

Mycoplasma hyopneumoniae Bacterin

Mycoplasma
Hyopneumoniae
Bacterin

Mycogard-1™ | 250 ML
250 DOSES



This product has been shown to be effective for the vaccination of healthy piglets, 10 days of age or older against Mycoplasma Hyopneumoniae. To review efficacy and safety data, go to productdata.aphis.usda.gov.

ADMINISTRATION AND DOSAGE: Dose: 1.0 ml intramuscularly (I.M.).

CAUTION: For animal use only. Do not mix with other products, except as specified on this label. In case of human exposure, contact a physician. Store at 2 to 8°C (35°F to 45°F). Do not freeze. Shake well before use. Use entire contents upon opening. Do not vaccinate within 21 days before slaughter. Transient reaction may occur at the injection site. If anaphylaxis occurs, use epinephrine or equivalent. Contains merthiolate (Thimerosal) as a preservative. This product has not been tested in pregnant animals. Revaccination interval not established.

Product No:
Serial No:
Exp. Date:

Manufactured by:
Pharmgate Biologics
2575 University Avenue
West, Suite 100,
St. Paul, MN 55114
612-256-0930
VLN 329 / PCN 2775.06

Visit USDA website for full label indications & technical information:
productdata.aphis.usda.gov

Pharmgate Animal Health swine vaccines/bacterins		
Product	Doses	Pharmgate Product Number
PRRSGard® (1 dose)	100	23010-100
PRRSGard® (1 dose)	250	23010-250
Mycogard® (2 dose)	50	20020-50
Mycogard® (2 dose)	250	20020-250
Mycogard-1™ (1 dose)	50	21010-50
Mycogard-1™ (1 dose)	250	21010-250
CircoGard® (1 dose)	50	22010-50
CircoGard® (1 dose)	250	22010-250
Circo/Mycogard® (1 dose)	50	25010-50
Circo/Mycogard® (1 dose)	250	25010-250