Porcine Circovirus Vaccine, Type 2, Killed Baculovirus Vector, Mycoplasma Hyopneumoniae Bacterin

Circo/MycoGard®

50 ML **50** DOSES



This product has been shown to be effective for the vaccination of healthy piglets, 10 days of age or older against Porcine Circovirus, Type 2b, and Mycoplasma Hyopneumoniae. The duration of immunity has not been established. This product is composed of PCV2b.

For more information regarding efficacy and safety data, see productdata.aphis.usda.gov.

ADMINISTRATION AND DOSAGE: Dose: 1.0 ml intramuscularly (I.M.).

CAUTION: For animal use only. Do not mix with other products, except as specified on this label. In case of human exposure, contact a physician. Store at 2 to 8°C (35°F to 45°F). Do not freeze. Shake well before use. Use entire contents upon opening. Do not vaccinate within 21 days before slaughter. Transient reaction may occur at the injection site. If anaphylaxis occurs, use epinephrine or equivalent. Contains merthiolate (Thimerosal) as a preservative. This product has not been tested in pregnant animals. For advice on revaccination frequency, consult your veterinarian.

1	
1	Т
1	ı
1	1
1	1
I .	1
1	1
	_
Draduat Na	

Product No:

Serial No:

Exp. Date:

Manufactured by:

Pharmgate Biologics Inc 2575 University Avenue West, Suite 100, St. Paul, MN 55114 612-256-0930

VLN 329 / PCN 49K5.R1