

Pennchlor[®] 100 Hi-Flo[™]

Chlortetracycline Type A Medicated Article

Active Drug Ingredients:		Chlortetracycline as Chlortetracycline Calcium Complex equivalent to 100 grams Chlortetracycline Hydrochloride / lb.	
Ingredients:		Chlortetracycline, Calcium Carbonate, Roughage Products and Mineral Oil.	
For Use in the Manufacture of Medicated Feeds CAUTION: For use in Dry Feeds ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS			
Directions For Use			
Indications For Use	Use Levels of Chlortetracycline	lbs. of Pennchlor 100 Hi-Flo [®] per ton	Indications For Use
Chickens			Swine
For Chickens: Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.	100-200g/ton	1.0-2.0	For Swine: Reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by <i>Group E Streptococci</i> susceptible to chlortetracycline.
For Chickens: Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.	200-400g/ton	2.0-4.0	For Breeding Swine: Control of leptospirosis (reducing the instances of abortions and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to chlortetracycline. (Feed continuously for not more than 14 days)
For Chickens: Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to chlortetracycline. (Feed for 5 days) WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. WITHDRAW 24 HOURS PRIOR TO SLAUGHTER.	500g/ton	5.0	For Swine: Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Feed for not more than 14 days) WARNING: ZERO-DAY WITHDRAWAL PERIOD.
Turkeys			CALVES, BEEF CATTLE, AND NONLACTATING DAIRY CATTLE WARNING: A WITHDRAWAL PERIOD HAS NOT BEEN ESTABLISHED FOR THIS PRODUCT IN PRE-RUMINATING CALVES. DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL.
For Turkeys: Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.	200g/ton	2.0	For Growing Cattle (over 400 lbs.): For the reduction of the incidence of liver abscesses. WARNING: ZERO-DAY WITHDRAWAL PERIOD.
For Turkeys: Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.	400g/ton	4.0	For Beef Cattle: For the control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella</i> spp. susceptible to chlortetracycline. WARNING: ZERO-DAY WITHDRAWAL PERIOD.
For Turkeys: Turkey poults not over 4 weeks of age: Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to chlortetracycline.	400g/ton	4.0	For Beef Cattle (under 700 lbs.): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline. WARNING: ZERO-DAY WITHDRAWAL PERIOD.
For Turkeys: Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.	25 mg/lb body weight/day		For Beef Cattle (over 700 lbs.): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline. WARNING: ZERO-DAY WITHDRAWAL PERIOD.
Sheep			For Calves, Beef, and Nonlactating Dairy Cattle: For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Treat for not more than 5 days) WARNING: ZERO-DAY WITHDRAWAL PERIOD.
For Breeding Sheep: Reducing the incidence of (vibronic) abortion caused by <i>Campylobacter fetus</i> infection susceptible to chlortetracycline. WARNING: ZERO-DAY WITHDRAWAL PERIOD.	80 mg/head/day		
Approved by FDA under NADA # 138-935 Pennchlor 100 Hi-Flo [®] is a registered trademark of Pharmgate Inc.			