

Pennchlor® 90 G™

Chlortetracycline Type A Medicated Article

Active Drug Ingredients:		Chlortetracycline as Chlortetracycline Calcium Complex equivalent to 90 grams Chlortetracycline Hydrochloride / lb.	
Ingredients:		Chlortetracycline, Calcium Carbonate, Roughage Products and Mineral Oil.	
For Use in the Manufacture of Medicated Feeds			
CAUTION: For use in Dry Feeds ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS			
Directions For Use			
Indications For Use		Indications For Use	
	Use Levels of Chlortetracycline	lbs. of Pennchlor 90 G® per ton	
<p>Chickens</p> <p>For Chickens: Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</p> <p style="text-align: center;">100-200g/ton 1.11-2.22</p> <p>For Chickens: Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</p> <p style="text-align: center;">200-400g/ton 2.22-4.44</p> <p>For Chickens: Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to chlortetracycline. (Feed for 5 days) WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. WITHDRAW 24 HOURS PRIOR TO SLAUGHTER.</p> <p style="text-align: center;">500g/ton 5.55</p>		<p>Swine</p> <p>For Swine: Reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by <i>Group E Streptococci</i> susceptible to chlortetracycline.</p> <p style="text-align: center;">50-100g/ton 0.555-1.11</p> <p>For Breeding Swine: Control of leptospirosis (reducing the instances of abortions and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to chlortetracycline. (Feed continuously for not more than 14 days)</p> <p style="text-align: center;">400g/ton 4.44</p> <p>For Swine: Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Feed for not more than 14 days) WARNING: ZERO-DAY WITHDRAWAL PERIOD.</p> <p style="text-align: center;">Feed approximately 400 g/t, varying with body weight and feed consumption to provide 10 mg/lb body weight/day</p>	
<p>Turkeys</p> <p>For Turkeys: Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</p> <p style="text-align: center;">200g/ton 2.22</p> <p>For Turkeys: Control of hexamitiasis caused by <i>Hexamita meleagridis</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</p> <p style="text-align: center;">400g/ton 4.44</p> <p>For Turkeys: Turkey poults not over 4 weeks of age: Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to chlortetracycline.</p> <p style="text-align: center;">400g/ton 4.44</p> <p>For Turkeys: Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</p> <p style="text-align: center;">25 mg/lb body weight/day</p>		<p>CALVES, BEEF CATTLE, AND NONLACTATING DAIRY CATTLE WARNING: A WITHDRAWAL PERIOD HAS NOT BEEN ESTABLISHED FOR THIS PRODUCT IN PRE-RUMINATING CALVES. DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL.</p> <p>For Growing Cattle (over 400 lbs.): For the reduction of the incidence of liver abscesses. WARNING: ZERO-DAY WITHDRAWAL PERIOD.</p> <p style="text-align: center;">70 mg/head/day</p> <p>For Beef Cattle: For the control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella</i> spp. susceptible to chlortetracycline. WARNING: ZERO-DAY WITHDRAWAL PERIOD.</p> <p style="text-align: center;">350 mg/head/day</p> <p>For Beef Cattle (under 700 lbs.): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline. WARNING: ZERO-DAY WITHDRAWAL PERIOD.</p> <p style="text-align: center;">350 mg/head/day</p> <p>For Beef Cattle (over 700 lbs.): Control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline. WARNING: ZERO-DAY WITHDRAWAL PERIOD.</p> <p style="text-align: center;">0.5 mg/lb body weight/day</p> <p>For Calves, Beef, and Nonlactating Dairy Cattle: For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Treat for not more than 5 days) WARNING: ZERO-DAY WITHDRAWAL PERIOD.</p> <p style="text-align: center;">10 mg/lb body weight/day</p>	
<p>Sheep</p> <p>For Breeding Sheep: Reducing the incidence of (vibriotic) abortion caused by <i>Campylobacter fetus</i> infection susceptible to chlortetracycline. WARNING: ZERO-DAY WITHDRAWAL PERIOD.</p> <p style="text-align: center;">80 mg/head/day</p>			
<p>Approved by FDA under NADA # 138-935 Pennchlor 90 G® is a registered trademark of Pharmgate Inc.</p>			