

NOTE: To achieve the approved combined indications for suppression of estrus (heat), control of coccidiosis caused by *Eimeria bovis* and *E. zuernii*, and for treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to chlortetracycline in replacement dairy heifers up to 800 pounds and less than 20 months of age and replacement beef heifers up to 800 pounds, the following Blue Bird labels must be followed, in which the melengestrol acetate Type C top-dress medicated feed is top dressed onto or mixed at feeding with a Type C medicated feed containing 500 to 4,000 g/ton of chlortetracycline and 30 to 181.8 g/ton of lasalocid for not more than 5 days.

Feeding of this combination requires a Veterinary Feed Directive.

**Replacement Dairy and Beef Heifer Feed – Top Dress
Melengestrol Acetate ES/Chlortetracycline/Lasalocid COCCI
Type C Medicated Feed
(melengestrol acetate Type C top-dress medicated feed)**

INDICATIONS FOR USE

For suppression of estrus (heat) in replacement dairy heifers up to 800 pounds and less than 20 months of age and replacement beef heifers up to 800 pounds.

ACTIVE DRUG INGREDIENT

Melengestrol acetate*.....0.5 to 2 g/ton (0.25 to 1 mg/lb)**

GUARANTEED ANALYSIS

Crude Protein (Min)	_____	%
NPN ¹ (Max)	_____	%
Crude Fat (Min)	_____	%
Crude Fiber (Max).....	_____	%
Calcium ¹ (Min).....	_____	%
Calcium ¹ (Max).....	_____	%
Phosphorus (Min).....	_____	%
Salt ¹ (Min).....	_____	%
Salt ¹ (Max)	_____	%
Sodium ² (Min).....	_____	%
Sodium ² (Max)	_____	%
Potassium ¹ (Min).....	_____	%
Vitamin A ¹ (Min)	_____	%

¹ Guaranteed required only when nutrient added except when feed is intended, represented, or serves as a principal source of the nutrient.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Must be top dressed onto or mixed at feeding with a Type C top-dress medicated feed applied onto or mix with a Type C medicated feed containing 500 to 4,000 g/ton of chlortetracycline and 30 to 181.8 g/ton lasalocid for not more than 5 days. After completing feeding this combination, continue feeding a Type C top-dress medicated feed containing melengestrol acetate alone for a total time not exceeding 24 days of feeding.

Feed 0.5 to 2 lb(s) per head per day (specify one level) of this Type C top-dress medicated feed to deliver 0.5 mg/head/day melengestrol acetate.

Examples in the table below may be used as a guide in determining the feeding rate based on desired mg melengestrol acetate per head per day and the concentration of the Type C top-dress medicated feed.

Amount of Type C top-dress medicated feed fed (lb/head/day)	Melengestrol acetate (mg/head/day)	Melengestrol acetate concentration in Type C top-dress medicated feed, g/ton (mg/lb)
0.5	0.5	2 (1)
2	0.5	0.5 (0.25)

WARNINGS



Withdrawal Periods

No withdrawal period is required when used according to labeling.



User Safety Warnings: Not for use in humans. Keep out of reach of children.

CAUTION

Not effective in steers and spayed heifers.

Do not exceed 24 days of feeding.

A reduced conception rate can be expected if heifers are bred at estruses observed within 1 to 12 days after withdrawal of melengestrol acetate, whereas heifers bred at subsequent observed estruses are expected to have normal conception rates.

QUESTIONS/COMMENTS

To report suspected adverse events or side effects, or for technical assistance contact Pharmgate Inc. at 1-800-380-6099 or www.pharmgate.com. For additional information about reporting side effects for animal drugs, contact FDA at 1-888- FDA-VETS or <http://www.fda.gov/reportanimalae>.

Lot Number (if applicable): _____
 Expiration dating not required.

Approved by FDA under ANADA # 200-776

MANUFACTURED BY

Blue Bird Feed Company
 Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

* MGA® is the proprietary name of melengestrol acetate Type A medicated article (NADA # 039-402).
 ** Final printed label on melengestrol acetate Type C top-dress medicated feed must bear a single concentration of the drug.

Pharmgate Inc. [9 July 2024]

Chlortetracycline/Lasalocid COCCI — CATTLE FEED
TYPE C Medicated Feed
(chlortetracycline and lasalocid Type C medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in cattle weighing up to 800 pounds.

ACTIVE DRUG INGREDIENTS*

Chlortetracycline^a.....500 to 4,000 g/ton**
Lasalocid^b (as lasalocid sodium).....30 to 181.8 g/ton**

GUARANTEED ANALYSIS

Crude Protein, not less than..... _____ %
(This includes not more than ___ % equivalent crude protein from non-protein nitrogen)¹
Crude Fat, not more than..... _____ %
Crude Fiber, not less than..... _____ %
Crude Fiber, not more than..... _____ %
Calcium, not less than _____ %
Calcium, not more than _____ %
Phosphorus, not less than _____ %
Potassium, not less than _____ %
Salt¹, not less than _____ %
Salt¹, not more than..... _____ %
Sodium², not less than..... _____ %
Sodium², not more than _____ %
Vitamin A¹, not less than..... _____ IU/lb

¹ If added

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

FEEDING DIRECTIONS

Hand feed continuously for not more than 5 days to provide 10 mg chlortetracycline per lb. body weight per day and 1 mg lasalocid per 2.2 lb. body weight per day to cattle weighing up to 800 lb. with a maximum of 360 mg of lasalocid per head per day. The following table provides examples of feeding rates and amounts:

Body Wt. Lb.	Feed Consumption		Chlortetracycline		Lasalocid	
	% Body Wt.	lb/head/day	Amount mg/lb BW/day	Feeding Rate g/ton	Amount mg/head/day	Feeding Rate g/ton
200	0.5	1	10	4,000	90.9	181.8
	1	2	10	2,000	90.9	90.9
	2	4	10	1,000	90.9	45.5
	3	6	10	667	90.9	30.3
800	0.5	4	10	4,000	360	180
	1	8	10	2,000	360	90
	2	16	10	1,000	360	45
	3	24	10	667	360	30

Caution

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

Warning: No withdrawal period is required. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Lot Number (if applicable): _____

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

*Final printed label on formulated Type C medicated feed must bear a single concentration for each drug.

**The ratio between chlortetracycline and lasalocid must be 22:1.

^aDeracin™ is the proprietary name of chlortetracycline Type A medicated article (ANADA 200-510).

^bBOVATEC® is the proprietary name of lasalocid sodium Type A medicated article (NADA 096-298).

Pharmgate Inc. [9 July 2024]