PRODUCT PROFILE

Tulissin®-100-
(tulathromycin injection)

Product Description

- Each mL of Tulissin 100 (tulathromycin injection) injectable solution contains 100 mg of tulathromycin.
- For use by or on the order of a licensed veterinarian.

Formulation

- 100 mg of tulathromycin / mL
- For use in swine*

Indication

- **Swine:** Tulissin 100 injectable solution is indicated for the treatment of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae, Pasteurella multocida, Bordetella bronchiseptica, Haemophilus parasuis* and *Mycoplasma hyopneumoniae*; and for the control of SRD associated with *Actinobacillus pleuropneumoniae, Pasteurella multocida* and *Mycoplasma hyopneumoniae* in groups of pigs where SRD has been diagnosed.

Packaging

- Tulissin 100 injectable solution is available in the following package sizes: 50 mL vial; 100 mL vial; 250 mL vial; 500 mL vial.

Dosage/Administration

- **Swine:** Inject intramuscularly as a single dose in the neck at a dosage of 2.5 mg/kg (0.25 mL/22 lb) body weight (BW). Do not inject more than 2.5 mL per injection site.

Precautions

- **Swine:** The effects of Tulissin 100 injectable solution on porcine reproductive performance, pregnancy, and lactation have not been determined. Intramuscular injection can cause a transient local tissue reaction that may result in trim loss of edible tissue at slaughter.

Withdrawal Period

- **Swine:** Swine intended for human consumption must not be slaughtered within 5 days from the last treatment.

Storage

- Store at or below 30° C (86° F). Use within 45 days of first puncture and puncture a maximum of 20 times. Consider using automatic injection equipment or a repeater syringe. When using a needle or draw-off spike larger than 16 gauge, discard any remaining product immediately after use.

Key Features

- Goes to work in minutes.[1]
- Concentrates in the most susceptible areas of the respiratory system.
- Provides nine days of lung activity to treat and control SRD.[2]
- New, patented[3] container design features an easy grip silicone shell that offers excellent shock-absorption properties. The innovative shell helps dramatically reduce the risk of economic loss, environmental contamination and lack of compliance if treatment is stopped.
- This exclusive packaging protects 250 mL and 500 mL vials against breakage, providing 92% resistance[4] when dropped multiple times from a height of up to four feet.
- Scan the QR code to see the protective shell in action.

IMPORTANT SAFETY INFORMATION TULISSIN® 100 (tulathromycin injection):
Ensure a pre-slaughter withdrawal time of five (5) days in swine. The effects of tulathromycin on swine reproductive performance, pregnancy and lactation have not been determined. Do not use in animals known to be hypersensitive to the product.

*Also approved for use in Beef and Non-Lactating Dairy Cattle, and Suckling, Dairy and Veal Calves. For more information, contact your Pharmgate Representative.

The labeling contains complete use information, including cautions and warnings. Always read, understand and follow the labeling and use directions. See the back page for use directions and additional information.

Pharmgate ANIMAL HEALTH

Tulissin®100

(tulathromycin injection)
Injectable Solution

Antibiotic
100 mg of tulathromycin/mL
For use in beef cattle (including suckling calves), non-lactating dairy cattle (including dairy calves), veal calves, and swine.
Not for use in female dairy cattle 20 months of age or older.

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION

TULISSIN 100 Injectable Solution is a ready-to-use sterile parenteral preparation containing tulathromycin, a semi-synthetic macrolide antibiotic of the subclass triamilide. Each mL of TULISSIN 100 contains 100 mg of tulathromycin, 500 mg propylene glycol, 19.2 mg citric acid and 5 mg monothioglycerol. Sodium hydroxide or hydrochloric acid may be added to adjust pH. TULISSIN 100 consists of an equilibrated mixture of two isomeric forms of tulathromycin in a 9:1 ratio. Structures of the isomers are shown below.
Figure 1.



The chemical names of the isomers are (2R,3S, 4R, 5R, 8R, 10R, 11R, 12S, 13S, 14R)-13 [[2,6-dideoxy-3-C-methyl-3-O-methyl-4-C-[(propylamino) methyl]-α-L-ribo-hexopyranosyl]oxy]-2-ethyl-3,4,10-trihydroxy-3, 5,8,10,12,14-hexamethyl-11-[[3,4,6-trideoxy-3-(dimethylamino)-β-D-xylo-hexopyranosyl]-oxy]-1-oxa-6-azacyclopentadecan-15-one and (2R, 3R,6R,8R,9R,10S,11S,12R)-11-[[2,6-dideoxy-3-C-methyl-3-O-methyl-4-C-[(propylamino)methyl]-α-L-ribo-hexopyranosyl] oxy]-2-[(1R,2R)-1,2-dihydroxy-1-methylbutyl]-8-hydroxy-3,6,8,10,12-pentamethyl-9-[[3,4,6-trideoxy-3-(dimethylamino)-β-D-xylo-hexopyranosyl]oxy]-1-oxa-4-azacyclotridecan-13-one, respectively.

INDICATIONS

Beef and Non-Lactating Dairy Cattle
BRD - TULISSIN 100 Injectable Solution is indicated for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica, Pasteurella multocida, Histophilus somni,* and *Mycoplasma bovis*; and for the control of respiratory disease in cattle at high risk of developing BRD associated with *Mannheimia haemolytica, Pasteurella multocida, Histophilus somni,* and *Mycoplasma bovis*.

IBK - TULISSIN 100 Injectable Solution is indicated for the treatment of infectious bovine keratoconjunctivitis (IBK) associated with *Moraxella bovis*.

Foot Rot - TULISSIN 100 Injectable Solution is indicated for the treatment of bovine foot rot (interdigital necrobacillosis) associated with *Fusobacterium necrophorum* and *Porphyromonas levii*.

Suckling Calves, Dairy Calves, and Veal Calves
BRD - TULISSIN 100 Injectable Solution is indicated for the treatment of BRD associated with *M. haemolytica, P. multocida, H. somni,* and *M. bovis*.

Swine
TULISSIN 100 Injectable Solution is indicated for the treatment of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae, Pasteurella multocida, Bordetella bronchiseptica, Haemophilus parasuis,* and *Mycoplasma hyopneumoniae*; and for the control of SRD associated with *Actinobacillus pleuropneumoniae, Pasteurella multocida,* and *Mycoplasma hyopneumoniae* in groups of pigs where SRD has been diagnosed.

DOSAGE AND ADMINISTRATION

Cattle
Inject subcutaneously as a single dose in the neck at a dosage of 2.5 mg/kg (1.1 mL/100 lb) body weight (BW). Do not inject more than 10 mL per injection site.
Table 1. TULISSIN 100 Injectable Solution Cattle Dosing Guide

Animal Weight (Pounds)	Dose Volume (mL)
100	1.1
200	2.3
300	3.4
400	4.5
500	5.7
600	6.8
700	8.0
800	9.1
900	10.2
1000	11.4

Swine
Inject intramuscularly as a single dose in the neck at a dosage of 2.5 mg/kg (0.25 mL/22 lb) BW. Do not inject more than 2.5 mL per injection site.
Table 2. TULISSIN 100 Injectable Solution Swine Dosing Guide

Animal Weight (Pounds)	Dose Volume (mL)
15	0.2
30	0.3
50	0.6
70	0.8
90	1.0
110	1.3
130	1.5
150	1.7
170	1.9
190	2.2
210	2.4
230	2.6
250	2.8
270	3.1
290	3.3

CONTRAINDICATIONS

The use of TULISSIN 100 Injectable Solution is contraindicated in animals previously found to be hypersensitive to the drug.

WARNINGS

FOR USE IN ANIMALS ONLY.
NOT FOR HUMAN USE.
KEEP OUT OF REACH OF CHILDREN.
NOT FOR USE IN CHICKENS OR TURKEYS.

RESIDUE WARNINGS

Cattle
Cattle intended for human consumption must not be slaughtered within 18 days from the last treatment. This drug is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows.

Swine
Swine intended for human consumption must not be slaughtered within 5 days from the last treatment.

PRECAUTIONS

Cattle
The effects of TULISSIN 100 Injectable Solution on bovine reproductive performance, pregnancy, and lactation have not been determined. Subcutaneous injection can cause a transient local tissue reaction that may result in trim loss of edible tissue at slaughter.

Swine
The effects of TULISSIN 100 Injectable Solution on porcine reproductive performance, pregnancy, and lactation have not been determined. Intramuscular injection can cause a transient local tissue reaction that may result in trim loss of edible tissue at slaughter.

ADVERSE REACTIONS

Cattle
In one BRD field study, two calves treated with tulathromycin injection at 2.5 mg/kg BW exhibited transient hypersalivation. One of these calves also exhibited transient dyspnea, which may have been related to pneumonia.

Swine
In one field study, one out of 40 pigs treated with tulathromycin injection at 2.5 mg/kg BW exhibited mild salivation that resolved in less than four hours.

POST APPROVAL EXPERIENCE

The following adverse events are based on post approval adverse drug experience reporting. Not all adverse events are reported to the FDA CVM. It is not always possible to reliably estimate the adverse event frequency or establish a causal relationship to product exposure using these data. The following adverse events are listed in decreasing order of reporting frequency in cattle: Injection site reactions and anaphylaxis/anaphylactoid reactions.
For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or http://www.fda.gov/reportanimalae.

CLINICAL PHARMACOLOGY

At physiological pH, tulathromycin (a weak base) is approximately 50 times more soluble in hydrophilic than hydrophobic media. This solubility profile is consistent with the extracellular pathogen activity typically associated with the macrolides.[1] Markedly higher tulathromycin concentrations are observed in the lungs as compared to the plasma. The extent to which lung concentrations represent free (active) drug was not examined. Therefore, the clinical relevance of these elevated lung concentrations is undetermined.
Although the relationship between tulathromycin and the characteristics of its antimicrobial effects has not been characterized, as a class, macrolides tend to be primarily bacteriostatic, but may be bactericidal against some pathogens.[2] They also tend to exhibit concentration independent killing; the rate of bacterial eradication does not change once serum drug concentrations reach 2 to 3 times the minimum inhibitory concentration (MIC) of the targeted pathogen. Under these conditions, the time that serum concentrations remain above the MIC becomes the major determinant of antimicrobial activity. Macrolides also exhibit a post-antibiotic effect (PAE), the duration of which tends to be both drug and pathogen dependent. In general, by increasing the macrolide concentration and the exposure time, the PAE will increase to some maximal duration. Of the two variables, concentration and exposure time, drug concentration tends to be the most powerful determinant of the duration of PAE.

Tulathromycin is eliminated from the body primarily unchanged via biliary excretion.
[1] *Carbon, C. 1998. Pharmacodynamics of Macrolides, Azalides, and Streptogramins: Effect on Extracellular Pathogens. Clin. Infect. Dis.,* ***27****:28-32.*
[2] *Nightingale, C.J. 1997. Pharmacokinetics and Pharmacodynamics of Newer Macrolides. Pediatr. Infect. Dis. J.,* ***16****:439-443.*

Cattle
Following subcutaneous administration into the neck of feeder calves at a dosage of 2.5 mg/kg BW, tulathromycin is rapidly and nearly completely absorbed. Peak plasma concentrations generally occur within 15 minutes after dosing and product relative bioavailability exceeds 90%. Total systemic clearance is approximately 170 mL/hr/kg. Tulathromycin distributes extensively into body tissues, as evidenced by volume of distribution values of approximately 11 L/kg in healthy ruminating calves.[3] This extensive volume of distribution is largely responsible for the long elimination half-life of this compound [approximately 2.75 days in the plasma (based on quantifiable terminal plasma drug concentrations) versus 8.75 days for total lung concentrations (based on data from healthy animals)]. Linear pharmacokinetics are observed with subcutaneous doses ranging from 1.27 mg/kg BW to 5.0 mg/kg BW. No pharmacokinetic differences are observed in castrated male versus female calves.
[3] *Clearance and volume estimates are based on intersubject comparisons of 2.5 mg/kg BW administered by either subcutaneous or intravenous injection.*

Swine
Following intramuscular administration to feeder pigs at a dosage of 2.5 mg/kg BW, tulathromycin is completely and rapidly absorbed (T_{max} ~0.25 hour). Subsequently, the drug rapidly distributes into body tissues, achieving a volume of distribution exceeding 15 L/kg. The free drug is rapidly cleared from the systemic circulation ($CL_{systemic}$ = 187 mL/hr/kg). However, it has a long terminal elimination half-life (60 to 90 hours) owing to its extensive volume of distribution. Although pulmonary tulathromycin concentrations are substantially higher than concentrations observed in the plasma, the clinical significance of these findings is undetermined. There are no gender differences in swine tulathromycin pharmacokinetics.

MICROBIOLOGY

Cattle
Tulathromycin has demonstrated *in vitro* activity against *Mannheimia haemolytica, Pasteurella multocida, Histophilus somni,* and *Mycoplasma bovis*, four pathogens associated with BRD; against *Moraxella bovis* associated with IBK; and against *Fusobacterium necrophorum* and *Porphyromonas levii* associated with bovine foot rot.
The MICs of tulathromycin against indicated BRD and IBK pathogens were determined using methods recommended by the Clinical and Laboratory Standards Institute (CLSI, M31-A2). The MICs against foot rot pathogens were also determined using methods recommended by the CLSI (M11-A6). All MIC values were determined using the 9:1 isomer ratio of this compound.

BRD - The MICs of tulathromycin were determined for BRD isolates obtained from calves enrolled in therapeutic and at-risk field studies in the U.S. in 1999. In the therapeutic studies, isolates were obtained from pre-treatment nasopharyngeal swabs from all study calves, and from lung swabs or lung tissue of saline-treated calves that died. In the at-risk studies, isolates were obtained from nasopharyngeal swabs of saline-treated non-responders, and from lung swabs or lung tissue of saline- treated calves that died. The results are shown in Table 3.

IBK - The MICs of tulathromycin were determined for *Moraxella bovis* isolates obtained from calves enrolled in IBK field studies in the U.S. in 2004. Isolates were obtained from pre-treatment conjunctival swabs of calves with clinical signs of IBK enrolled in the tulathromycin injection and saline-treated groups. The results are shown in Table 3.

Foot Rot - The MICs of tulathromycin injection were determined for *Fusobacterium necrophorum* and *Porphyromonas levii* obtained from cattle enrolled in foot rot field studies in the U.S. and Canada in 2007. Isolates were obtained from pre-treatment interdigital biopsies and swabs of cattle with clinical signs of foot rot enrolled in the tulathromycin injection and saline-treated groups. The results are shown in Table 3.

Table 3. Tulathromycin minimum inhibitory concentration (MIC) values* for indicated pathogens isolated from field studies evaluating BRD and IBK in the U.S. and from foot rot field studies in the U.S. and Canada.

Indicated Pathogen	Date isolated	No. of isolates	MIC_{50} † (µg/mL)	MIC_{90} † (µg/mL)	MICrange (µg/mL)
Mannheimia haemolytica	1999	642	2	2	0.5 to 64
Pasteurella multocida	1999	221	0.5	1	0.25 to 64
Histophilus somni	1999	36	4	4	1 to 4
Mycoplasma bovis	1999	43	0.125	1	≤ 0.063 to > 64
Moraxella bovis	2004	55	0.5	0.5	0.25 to 1
Fusobacterium necrophorum	2007	116	2	64	≤ 0.25 to > 128
Porphyromonas levii	2007	103	8	128	≤ 0.25 to > 128

* The correlation between *in vitro* susceptibility data and clinical effectiveness is unknown.
† The lowest MIC to encompass 50% and 90% of the most susceptible isolates, respectively.

Swine
In vitro activity of tulathromycin has been demonstrated against *Actinobacillus pleuropneumoniae, Pasteurella multocida, Bordetella bronchiseptica, Haemophilus parasuis,* and *Mycoplasma hyopneumoniae*.
The MICs of tulathromycin against indicated SRD pathogens were determined using methods recommended by the Clinical and Laboratory Standards Institute (CLSI, M31-A and M31-A3). MICs for *Haemophilus parasuis* were determined using Veterinary Fastidious Medium and were incubated up to 48 hours at 35 to 37°C in a CO_2-enriched atmosphere. All MIC values were determined using the 9:1 isomer ratio of this compound. Isolates obtained in 2000 and 2002 were from lung samples from saline-treated pigs and non-treated sentinel pigs enrolled in Treatment of SRD field studies in the U.S. and Canada. The results are shown in Table 4. Isolates obtained in 2007 and 2008 were from lung samples from saline-treated and tulathromycin injection-treated pigs enrolled in the Control of SRD field study in the U.S. and Canada. The results are shown in Table 4.

Table 4. Tulathromycin minimum inhibitory concentration (MIC) values* for indicated pathogens isolated from field studies evaluating SRD in the U.S. and Canada.

Indicated Pathogen	Date isolated	No. of isolates	MIC_{50} † (µg/mL)	MIC_{90} † (µg/mL)	MICrange (µg/mL)
Actinobacillus pleuropneumoniae	2000-2002	135	16	32	16 to 32
	2007-2008	88	16	16	4 to 32
Haemophilus parasuis	2000-2002	31	1	2	0.25 to > 64
Pasteurella multocida	2000-2002	55	1	2	0.5 to > 64
	2007-2008	40	1	2	≤ 0.03 to 2
Bordetella bronchiseptica	2000-2002	42	4	8	2 to 8

* The correlation between in vitro susceptibility data and clinical effectiveness is unknown.
† The lowest MIC to encompass 50% and 90% of the most susceptible isolates, respectively.

EFFECTIVENESS

Cattle
BRD – In a multi-location field study, 314 calves with naturally occurring BRD were treated with tulathromycin injection. Responses to treatment were compared to saline-treated controls. A cure was defined as a calf with normal attitude/activity, normal respiration, and a rectal temperature of ≤ 104°F on Day 14. The cure rate was significantly higher (P ≤ 0.05) in tulathromycin injection-treated calves (78%) compared to saline-treated calves (24%). There were two BRD-related deaths in the tulathromycin injection-treated calves compared to nine BRD-related deaths in the saline-treated calves.
Fifty-two tulathromycin injection-treated calves and 27 saline-treated calves from the multi-location field BRD treatment study had *Mycoplasma bovis* identified in cultures from pre-treatment nasopharyngeal swabs.
Of the 52 tulathromycin injection-treated calves, 37 (71.2%) calves were categorized as cures and 15 (28.8%) calves were categorized as treatment failures. Of the 27 saline-treated calves, 4 (14.8%) calves were categorized as cures and 23 (85.2%) calves were treatment failures.
A Bayesian meta-analysis was conducted to compare the BRD treatment success rate in young calves (calves weighing 250 lbs or less and fed primarily a milk-based diet) treated with tulathromycin injection to the success rate in older calves (calves weighing more than 250 lbs and fed primarily a roughage and grain-based diet) treated with tulathromycin injection. The analysis included data from four BRD treatment effectiveness studies conducted for the approval of tulathromycin injection in the U.S. and nine contemporaneous studies conducted in Europe. The analysis showed that the BRD treatment success rate in young calves was at least as good as the BRD treatment success rate in older calves. As a result, tulathromycin injection is considered effective for the treatment of BRD associated with *M. haemolytica, P. multocida, H. somni,* and *M. bovis* in suckling calves, dairy calves, and veal calves.
In another multi-location field study with 399 calves at high risk of developing BRD, administration of tulathromycin injection resulted in a significantly reduced incidence of BRD (11%) compared to saline-treated calves (59%). Effectiveness evaluation was based on scored clinical signs of normal attitude/activity, normal respiration, and a rectal temperature of ≤ 104°F on Day 14. There were no BRD-related deaths in the tulathromycin injection-treated calves compared to two BRD-related deaths in the saline-treated calves. Fifty saline-treated calves classified as non-responders in this study had *Mycoplasma bovis* identified in cultures of post-treatment nasopharyngeal swabs or lung tissue.
Two induced infection model studies were conducted to confirm the effectiveness of tulathromycin injection against *Mycoplasma bovis*. A total of 166 calves were inoculated intratracheally with field strains of *Mycoplasma bovis*. When calves became pyrexic and had abnormal respiration scores, they were treated with either tulathromycin injection (2.5 mg/kg BW) subcutaneously or an equivalent volume of saline. Calves were observed for signs of BRD for 14 days post-treatment, then were euthanized and necropsied. In both studies, mean lung lesion percentages were statistically significantly lower in the tulathromycin injection-treated calves compared with saline-treated calves (11.3% vs. 28.9%, P = 0.0001 and 15.0% vs. 30.7%, P < 0.0001).
IBK – Two field studies were conducted evaluating tulathromycin injection for the treatment of IBK associated with *Moraxella bovis* in 200 naturally-infected calves. The primary clinical endpoint of these studies was cure rate, defined as a calf with no clinical signs of IBK and no corneal ulcer, assessed on Days 5, 9, 13, 17, and 21. Time to improvement, defined as the first day on which a calf had no clinical signs of IBK in both eyes, provided that those scores were maintained at the next day of observation, was assessed as a secondary variable. At all time points, in both studies, the cure rate was significantly higher (P < 0.05) for tulathromycin injection-treated calves compared to saline-treated calves. Additionally, time to improvement was significantly less (P < 0.0001) in both studies for tulathromycin injection-treated calves compared to saline-treated calves.
Foot Rot - The effectiveness of tulathromycin injection for the treatment of bovine foot rot was evaluated in 170 cattle in two field studies. Cattle diagnosed with bovine foot rot were enrolled and treated with a single subcutaneous dose of tulathromycin injection (2.5 mg/kg BW) or an equivalent volume of saline. Cattle were clinically evaluated 7 days after treatment for treatment success, which was based on defined decreases in lesion, swelling, and lameness scores. In both studies, the treatment success percentage was statistically significantly higher in tulathromycin injection-treated calves compared with saline-treated calves (60% vs. 8%, P < 0.0001 and 83.3% vs. 50%, P = 0.0088).

Swine
In a multi-location field study to evaluate the treatment of naturally occurring SRD, 266 pigs were treated with tulathromycin injection. Responses to treatment were compared to saline-treated controls. Success was defined as a pig with normal attitude, normal respiration, and rectal temperature of < 104°F on Day 7. The treatment success rate was significantly greater (P ≤ 0.05) in tulathromycin injection-treated pigs (70.5%) compared to saline-treated pigs (46.1%).
M. hyopneumoniae was isolated from 106 saline-treated and non-treated sentinel pigs in this study.
Two induced infection model studies were conducted to confirm the effectiveness of tulathromycin injection against *M. hyopneumoniae*. Ten days after inoculation intranasally and intratracheally with a field strain of *M. hyopneumoniae*, 144 pigs were treated with either tulathromycin injection (2.5 mg/kg BW) intramuscularly or an equivalent volume of saline. Pigs were euthanized and necropsied 10 days post-treatment. The mean percentage of gross pneumonic lung lesions was statistically significantly lower (P < 0.0001) for tulathromycin injection-treated pigs than for saline-treated pigs in both studies (8.52% vs. 23.62% and 11.31% vs. 26.42%).
The effectiveness of tulathromycin injection for the control of SRD was evaluated in a multi-location natural infection field study. When at least 15% of the study candidates showed clinical signs of SRD, all pigs were enrolled and treated with tulathromycin injection (226 pigs) or saline (227 pigs). Responses to treatment were evaluated on Day 7. Success was defined as a pig with normal attitude, normal respiration, and rectal temperature of < 104°F. The treatment success rate was significantly greater (P < 0.05) in tulathromycin injection-treated pigs compared to saline-treated pigs (59.2% vs. 41.2%).

ANIMAL SAFETY

Cattle
Safety studies were conducted in feeder calves receiving a single subcutaneous dose of 25 mg/kg BW, or 3 weekly subcutaneous doses of 2.5, 7.5, or 12.5 mg/kg BW. In all groups, transient indications of pain after injection were seen, including head shaking and pawing at the ground. Injection site swelling, discoloration of the subcutaneous tissues at the injection site and corresponding histopathologic changes were seen in animals in all dosage groups.These lesions showed signs of resolving over time. No other drug-related lesions were observed macroscopically or microscopically.
An exploratory study was conducted in feeder calves receiving a single subcutaneous dose of 10, 12.5, or 15 mg/kg BW. Macroscopically, no lesions were observed. Microscopically, minimal to mild myocardial degeneration was seen in one of six calves administered 12.5 mg/kg BW and two of six calves administered 15 mg/kg BW.
A safety study was conducted in preruminant calves 13 to 27 days of age receiving 2.5 mg/kg BW or 7.5 mg/kg BW once subcutaneously. With the exception of minimal to mild injection site reactions, no drug-related clinical signs or other lesions were observed macroscopically or microscopically.

Swine
Safety studies were conducted in pigs receiving a single intramuscular dose of 25 mg/kg BW, or 3 weekly intramuscular doses of 2.5, 7.5, or 12.5 mg/kg BW. In all groups, transient indications of pain after injection were seen, including restlessness and excessive vocalization. Tremors occurred briefly in one animal receiving 7.5 mg/kg BW. Discoloration and edema of injection site tissues and corresponding histopathologic changes were seen in animals at all dosages and resolved over time. No other drug-related lesions were observed macroscopically or microscopically.

STORAGE CONDITIONS

Store at or below 30°C (86°F). Use within 45 days of first puncture and puncture a maximum of 20 times. Consider using automatic injection equipment or a repeater syringe. When using a needle or draw-off spike larger than 16 gauge, discard any remaining product immediately after use.

HOW SUPPLIED: TULISSIN 100 Injectable Solution is available in the following package sizes:
50 mL vial ; 100 mL vial ; 250 mL vial ; 500 mL vial

Manufactured for: Virbac AH, Inc. - P.O. Box 162059, Fort Worth, TX 76161 - Made in France

Approved by FDA under ANADA # 200-669

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Virbac AH, Inc. at 1-800-338-3659 or us.virbac.com. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or http://www.fda.gov/reportanimalae.

©2021 Virbac Corporation. All Rights Reserved.
TULISSIN is a registered trademark of Virbac S.A.
Rev: 06/22

TAKE TIME OBSERVE LABEL DIRECTIONS

1 Villarino N, Brown SA, Martín-Jiménez T. Understanding the pharmacokinetics of tulathromycin: a pulmonary perspective. *J Vet Pharmacol Ther.* 2014;37(3):211-221.

2 Waag TA, Bradford JR, Lucas MJ, et al. Duration of effectiveness of tulathromycin injectable solution in an *Actinobacillus pleuropneumoniae* respiratory-disease challenge model in swine. *J Swine Health Prod.* 2008;16(3):126–130.

3 Patent application n°WO2019201812 Registered international design n°DM/103483.

4 Internal study.

Marketed by

Pharmgate ANIMAL HEALTH

©2025 Virbac Corporation. All Rights Reserved. TULISSIN is a registered trademark of Virbac S.A. 003TUL-0825

Virbac Corporation 1301 Solana Boulevard, Building 2, Suite 2400
Westlake, TX 76262. https://vet-us.virbac.com/home.html