

Deracin™ (chlortetracycline) Veterinary Feed Directive for use in Psittacine Birds

Client: _____ Veterinarian: _____
Business or _____
Home _____
Address: _____ Address: _____
_____ Address: _____
Phone #: _____ Phone #: _____

Approximate number of **Psittacine Birds** to be treated: _____

Location of animals: _____

Special Instructions and/or other animal identifications:

Indication, Drug Level, and Duration of Use:

Psittacine Birds: Treatment of psittacine birds (parrots, macaws, cockatoos) suspected or known to be infected with psittacosis caused by *Chlamydia psittaci* sensitive to chlortetracycline.

Drug level: 10 mg/g of feed**Duration of use:** 45 days

Each bird should consume an amount of medicated feed equal to one-fifth of its body weight daily. During treatment, parrots, macaws, and cockatoos should be kept individually or in pairs in clean cages.

USE OF FEED CONTAINING THIS VETERINARY FEED DIRECTIVE (VFD) DRUG IN A MANNER OTHER THAN AS DIRECTED ON THE LABELING (EXTRA-LABEL USE) IS NOT PERMITTED.

Affirmation of Intent (for combination VFD drugs):

This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.

Warning: Psittacosis, avian chlamydiosis, or ornithosis is a reportable communicable disease, transmissible between wild animals and domestic birds, other animals and man. Contact appropriate public health and regulatory officials.

Caution: Aspergilliosis may occur following prolonged treatment.

VFD Issuance Date: _____

VFD Expiration Date: _____
Month/Day/Year
(Not to exceed 6 months from issuance date)

Veterinarian's signature: _____

Approved by FDA under ANADA # 200-510

Rev. July 29, 2025

Copy - Supplier

Copy – Client

Copy – Feed Mill