Chlortetracycline Swine Supplement
(chlortetracycline Type B medicated feed)

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated swine feed.

ACTIVE DRUG INGREDIENT

Chlortetracycline[a] ..... 2 – 40 g/lb*

GUARANTEED ANALYSIS

Crude Protein (Min)..... _____%
Lysine (Min)..... _____%
Crude Fat (Min)..... _____%
Crude Fiber (Max)..... _____%
Calcium (Min) ..... _____%
Calcium (Max)..... _____%
Phosphorus (Min) ..... _____%
Salt[1] (Min) ..... _____%
Salt[1] (Max)..... _____%
Sodium[2] (Min) ..... _____%
Sodium[2] (Max) ..... _____%
Selenium (Min)..... _____ppm

[1] Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.
[2] Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Blend with grains and non-medicated feed ingredients to provide a complete feed with the following grams of chlortetracycline per ton of complete feed.

Indications for Use	Use of Levels of Chlortetracycline
SWINE	
For Swine: Mix 25 to 50 pounds of Type B (2 g/lb) or 1 ¼ to 2 ½ pounds of Type B (40 g/lb) with non-medicated feed ingredients to make a ton of swine feed for reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by group E streptococci susceptible to chlortetracycline containing 50-100 g/ton to be fed continuously.	50-100 g/ton
For Breeding Swine: Mix 200 pounds of Type B (2 g/lb) or 10 pounds of Type B (40 g/lb) with non-medicated feed ingredients to make a ton of swine feed for the control of leptospirosis (reducing the incidence of abortions and shedding of leptospirae) caused by *Leptospira pomona* susceptible to chlortetracycline containing 400 g/ton to be feed continuously for not more than 14 days.	400 g/ton

For Swine: Mix 200 to 1,000 pounds of Type B (2 g/lb) or 10 to 50 pounds of Type B (40 g/lb) to provide 400 g/ton varying with body weight and feed consumption to provide 10 mg/lb body weight/day in the resulting Type C medicated feed with non-medicated feed ingredients to make a ton of swine feed for the treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella* Choleraesuis and bacterial pneumonia caused by *Pasteurella multocida* susceptible to chlortetracycline to be fed for not more than 14 days.	400 – 2,000 g/ton to provide 10 mg/lb body weight/day

Withdrawal Periods: Zero-day withdrawal period.

*The final printed label must bear a single drug concentration.

Net Weight_____lb (_____kg)
Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000

Lot Number: _____

[a]Sourced from Pennchlor® (chlortetracycline Type A medicated article); NADA # 138-935.

Pharmgate [Revision Date: September 2, 2025]