CTC / MONENSIN 0.5mg/ANA FE – BEEF FEEDLOT CATTLE FEED
(chlortetracycline and monensin Type C medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Indications for Use

For control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline and for improved feed efficiency in growing beef steers and heifers fed in confinement for slaughter over 700 lbs.

Active Drug Ingredients

Chlortetracycline[a] (as chlortetracycline calcium complex) equivalent to chlortetracycline hydrochloride……..... 33.33 to 100 g/ton*

Monensin, USP[b].....…...……..…… 5 to 40 g/ton*

Guaranteed Analysis

Crude protein (min)	.....	_____	%
NPN[1] (max)	.....	_____	%
Crude fat (min)	.....	_____	%
Crude fiber (max)	.....	_____	%
Calcium[1] (min)	.....	_____	%
Calcium[1] (max)	.....	_____	%
Phosphorus[1] (min)	.....	_____	%
Salt[1] (min)	.....	_____	%
Salt[1] (max)	.....	_____	%
Sodium[2] (min)	.....	_____	%
Sodium[2] (max)	.....	_____	%
Potassium[1] (min)	.....	_____	%
Vitamin A[1] (min)	.....	_____	IU/lb

[1] Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of nutrient.
[2]Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient as named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

Feeding Directions

Feed as the sole ration to provide 0.5 mg chlortetracycline per lb. body weight per day and 50 to 480 mg monensin per head per day. No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 grams per ton (360 mg monensin per head per day).

The following table provides examples of feeding rates and amount of drug per ton of feed:

Body Weight (lbs)	Feed Consumption (lbs per day)	Chlortetracycline in medicated feed		Monensin in medicated feed	
		mg/lb BW/day	g/ton	mg/head/day	g/ton
800	8	0.5	100	160	40
1000	15	0.5	66.67	225	30
1200	24	0.5	50	120	10
1400	42	0.5	33.33	105	5

Caution

For use in dry feeds only. Not for use in liquid feed supplements. Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning

Withdrawal Periods and Residue Warnings:
No withdrawal period is required when used according to labeling.
A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

User Safety Warnings:
Keep this and all drugs out of the reach of children. Not for human use.

Lot No. _____

Approved by FDA under NADA # 141-564

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000

NET WEIGHT ON BAG OR BULK

---

*Final printed label on formulated Type C medicated feed must bear a single concentration for each drug.
[a] Sourced from Pennchlor®, NADA # 138-935
[b] Sourced from Rumensin™, NADA # 95-735