

**Oxytetracycline Calf Supplement
Type B Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated calf feed for use in calf milk replacers or starter feeds.

ACTIVE DRUG INGREDIENT

Oxytetracycline hydrochloride. 2-20 g/lb*

GUARANTEED ANALYSIS

Crude Protein, (Min)	XX.XX %
NPN, (Max) ¹	X.XX%
Crude Fat, (Min)	X.XX%
Crude Fiber, (Max)	X.XX%
Calcium, (Min) ¹	X.XX%
Calcium, (Max) ¹	X.XX%
Phosphorus, (Min) ¹	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%
Potassium, (Min) ¹	X.XX%
Vitamin A, (Min) ³	___IU/Lb

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

³Other than precursors of Vitamin A, if added

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below. To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing as a complete feed.

Indications for Use

Oxytetracycline in
Type C medicated feed

For Calves: Mix an amount of Type B feed to provide approximately 10 mg/lb body weight/day varying with body weight and feed consumption with non-medicated feed ingredients to make the daily complete feed for treatment of bacterial enteritis caused by Escherichia coli susceptible to oxytetracycline. Feed continuously for 7-14 days as the sole ration.

10 mg/lb body weight/day

RESIDUE WARNING: 5-day withdrawal period at 10 mg/lb dosage.

*The medicated feed label must state a single drug concentration.

Net Weight _____ lb (_____ kg)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000