

**Oxytetracycline Broiler Chicken Ration for Reduction of Mortality Due to Air Sacculitis  
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

For reduction of mortality due to air sacculitis (air-sac-infection) caused by Escherichia coli susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to 500 grams oxytetracycline hydrochloride/ton.

GUARANTEED ANALYSIS

Crude Protein, (Min) . . . . .	XX.XX %
Lysine, (Min) . . . . .	X.XX%
Methionine, (Min) . . . . .	X.XX%
Crude Fat, (Min) . . . . .	X.XX%
Crude Fiber, (Max) . . . . .	X.XX%
Calcium, (Min) . . . . .	X.XX%
Calcium, (Max) . . . . .	X.XX%
Phosphorus, (Min) . . . . .	X.XX%
Salt, (Min) <sup>1</sup> . . . . .	X.XX%
Salt, (Max) <sup>1</sup> . . . . .	X.XX%
Sodium, (Min) <sup>2</sup> . . . . .	X.XX%
Sodium, (Max) <sup>2</sup> . . . . .	X.XX%

<sup>1</sup>If added

<sup>2</sup>Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO

DIRECTIONS FOR USE

Feed continuously 5 days as the sole ration.

**RESIDUE WARNING:** Do not feed to chickens producing eggs for human consumption. 24 hour withdrawal period. Do not use in low-calcium feed containing less than 0.55% dietary calcium. Use in such feeds may result in violative residues.

Net Weight \_\_\_\_\_ lb (\_\_\_\_\_ kg)  
Manufactured by  
Blue Bird Feed Company  
Blue Bird, MD 00000