# Oxytetracycline Cattle Ration for the Prevention and Treatment of the Early Stages of the Shipping Fever Complex Type C Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the prevention and treatment of the early stages of the shipping fever complex.

## ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to	grams* oxytetracycline hydrochloride/ton
to provide approximately 0.5-2.0 g/head/day varying with body weight and feed	consumption.

### **GUARANTEED ANALYSIS**

Crude Protein, (Min)	XX.XX %
NPN, (Max) <sup>1</sup>	X.XX%
Crude Fat, (Min)	X.XX%
Crude Fiber, (Max)	X.XX%
Calcium, (Min) <sup>1</sup>	X.XX%
Calcium, (Max) <sup>1</sup>	X.XX%
Phosphorus, (Min) <sup>1</sup>	X.XX%
Salt, (Min) <sup>1</sup>	X.XX%
Salt, (Max) <sup>1</sup>	X.XX%
Sodium, (Min) <sup>2</sup>	X.XX%
Sodium, (Max) <sup>2</sup>	X.XX%
Potassium, (Min) <sup>1</sup>	X.XX%
Vitamin A, $(Min)^3$	IU/Lb

<sup>&</sup>lt;sup>1</sup>If added

#### **INGREDIENTS**

Ingredients as defined by AAFCO

## DIRECTIONS FOR USE

Feed 3-5 days before and after arrival in feedlots.

RESIDUE WARNING: Zero-day withdrawal period.

\*The medicated feed label must state a single drug concentration.

Net Weight \_\_\_\_\_ lb (\_\_\_\_\_ kg)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000

<sup>&</sup>lt;sup>2</sup>Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

<sup>&</sup>lt;sup>3</sup>Other than precursors of Vitamin A, if added