

**Oxytetracycline Chicken Ration for Control of Infectious Synovitis and Control of Fowl Cholera  
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

For control of infectious synovitis caused by Mycoplasma synoviae and control of fowl cholera caused by Pasteurella multocida susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to \*100-200 grams oxytetracycline hydrochloride/ton.

GUARANTEED ANALYSIS

Crude Protein, (Min) . . . . .	XX.XX %
Lysine, (Min) . . . . .	X.XX%
Methionine, (Min) . . . . .	X.XX%
Crude Fat, (Min) . . . . .	X.XX%
Crude Fiber, (Max) . . . . .	X.XX%
Calcium, (Min) . . . . .	X.XX%
Calcium, (Max) . . . . .	X.XX%
Phosphorus, (Min) . . . . .	X.XX%
Salt, (Min) <sup>1</sup> . . . . .	X.XX%
Salt, (Max) <sup>1</sup> . . . . .	X.XX%
Sodium, (Min) <sup>2</sup> . . . . .	X.XX%
Sodium, (Max) <sup>2</sup> . . . . .	X.XX%

<sup>1</sup>If added

<sup>2</sup>Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO

DIRECTIONS FOR USE

Feed continuously for 7-14 days as the sole ration.

**RESIDUE WARNING:** Do not feed to chickens producing eggs for human consumption. Zero-day withdrawal period. Do not use in low-calcium feed containing less than 0.55% dietary calcium. Use in such feeds may result in violative residues.

\*The medicated feed label must state a single drug concentration.

Net Weight \_\_\_\_\_ lb (\_\_\_\_\_ kg)  
Manufactured by  
Blue Bird Feed Company

Blue Bird, MD 00000