Oxytetracycline Swine Ration for Treatment of Bacterial Enteritis and Bacterial Pneumonia Type C Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For treatment of bacterial enteritis caused by <u>Escherichia coli</u> and <u>Salmonella choleraeuis</u> and bacterial pneumonia caused by Pasteurella multocida susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to	grams* oxytetracycline
hydrochloride/ton varying with body weight and feed consumption to prov	vide approximately 10 mg/lb body
weight/day.	

GUARANTEED ANALYSIS

Crude Protein, (Min)	XX.XX %
Lysine, (Min)	X.XX%
Crude Fat, (Min)	X.XX%
Crude Fiber, (Max)	X.XX%
Calcium, (Min)	X.XX%
Calcium, (Max)	X.XX%
Phosphorus, (Min)	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, $(Min)^2$	X.XX%
Sodium, $(Max)^2$	X.XX%
Selenium, (Min)	ppm
Zinc, (Min)	ppm

¹If added

INGREDIENTS

Ingredients as defined by AAFCO

DIRECTIONS FOR USE

Feed continuously for 7-14 days as the sole ration.

RESIDUE WARNING: Zero-day withdrawal period.

*The medicated feed label must state a single drug concentration.

Net Weight ______ lb (_____ kg)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee