

NEO-OXY (neomycin sulfate, oxytetracycline) Veterinary Feed Directive for use in Turkeys

Client: _____	Veterinarian: _____
Business or Home Address: _____	Address: _____
Phone #: _____	Phone #: _____

Approximate number of turkeys to be treated: _____

Location of animals: _____

Special Instructions and/or other animal identifications:

Indication, Drug Level in Medicated Feed, and Duration of Use (check one):

- A)** For control of hexamitiasis caused by Hexamita meleagridis susceptible to oxytetracycline.
 Drug level: 100 g/ton
 Duration of use: _____ days (specify number of days from 7 to 14 days)

- B)** For control of infectious synovitis caused by Mycoplasma synoviae susceptible to oxytetracycline.
 Drug level: 200 g/ton
 Duration of use: _____ days (specify number of days from 7 to 14 days)

- C)** For control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline.
 Drug level: _____ g/ton in order to provide 25 mg / lb body weight / day
 Duration of use: _____ days (specify number of days from 7 to 14 days)

Caution: Use of feed containing this Veterinary Feed Directive (VFD) drug in a manner other than as directed on the labeling (extra-label use) is not permitted.

For use in Dry Feeds Only. Not for Use in Liquid Medicated Feeds.

▶

Residue Warnings:
Indication A: Do not feed to turkeys producing eggs for human consumption. Zero-day withdrawal period.
Indication B and C: Do not feed to turkeys producing eggs for human consumption. Withdraw 5 days before slaughter.

◀

Combination Use:

This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.

VFD Issuance Date: _____

VFD Expiration Date: _____
Month/Day/Year
 (Not to exceed 6 months from issuance date)

Veterinarian's signature: _____

Copy – Supplier

Copy – Client

Original – Veterinarian