Chlortetracycline Beef Cattle Feed – DERA[a] – BE-BP – TD
(chlortetracycline Type B medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS FOR USE

For calves, beef and non-lactating dairy cattle: Treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to chlortetracycline.

ACTIVE DRUG INGREDIENT

Chlortetracycline[a] ..... 20,001 – 80,000 g/ton*

GUARANTEED ANALYSIS

Crude Protein (Min) ..... _____%
NPN[1] (Max) ..... _____%
Crude Fat (Min) ..... _____%
Crude Fiber (Max) ..... _____%
Calcium[1] (Min) ..... _____%
Calcium[1] (Max) ..... _____%
Phosphorus[1] (Min) ..... _____%
Salt[1] (Min) ..... _____%
Salt[1] (Max) ..... _____%
Sodium[2] (Min) ..... _____%
Sodium[2] (Max) ..... _____%
Potassium[1] (Min) ..... _____%
Vitamin A[1] ..... _____IU/lb

[1] Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.
[2] Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by Association of American Feed Control Official (AAFCO).

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C top-dress medicated feed for cattle containing 4,000 – 20,000 g chlortetracycline per ton.

The following table provides examples of mixing rates:

Type B CTC concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C CTC concentration (g/ton)
40,000 (20 g/lb)	200	1,800	4,000
80,000 (40 g/lb)	100	1,900	4,000
80,000 (40 g/lb)	500	1,500	20,000

The resulting Type C medicated feed should be fed for not more than 5 days to provide chlortetracycline at the rate of 10 mg/pound of body weight per day. The medicated feed should be administered as a top dress or mixed into the daily ration. If top-dressed, spread uniformly on top of ration and allow sufficient space so that all cattle can eat at the same time.

WITHDRAWAL PERIODS AND RESIDUE WARNINGS

No withdrawal period is required when used according to labeling. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

USER SAFETY WARNINGS

Not for use in humans. Keep out of reach of children.

QUESTIONS/COMMENTS?

For technical assistance or to obtain a Safety Data Sheet(s), contact Pharmgate Animal Health LLC at 1-800-380-6099 or www.pharmgate.com. To report side effects, contact Pharmgate Animal Health LLC at 1-833-531-0114. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or http://www.fda.gov/reportanimalae.

Approved by FDA under ANADA # 200-510

Lot Number: _____

MANUFACTURED BY

Blue Bird Feed Mill
City, State, Zip

NET WEIGHT ON BAG OR BULK

*The final printed medicated feed label must state a single drug concentration.

[a] Sourced from Deracin™ (chlortetracycline Type A medicated article), ANADA # 200-510.

Pharmgate Inc. [29 July 2025]