Chlortetracycline Turkey Poult Feed – DERA[a]
(chlortetracycline Type B medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS FOR USE

For reduction of mortality due to paratyphoid caused by *Salmonella typhimurium* susceptible to chlortetracycline in turkey poults not over 4 weeks of age.

ACTIVE DRUG INGREDIENT

Chlortetracycline[a] ..... 401 – 80,000 g/ton*

GUARANTEED ANALYSIS

Crude Protein (Min) ..... _____%
Lysine (Min) ..... _____%
Methionine (Min) ..... _____%
Crude Fat (Min) ..... _____%
Crude Fiber (Max) ..... _____%
Calcium (Min) ..... _____%
Calcium (Max) ..... _____%
Phosphorus (Min) ..... _____%
Salt[1] (Min) ..... _____%
Salt[1] (Max) ..... _____%
Sodium[2] (Min) ..... _____%
Sodium[2] (Max) ..... _____%

[1] Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.
[2] Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by Association of American Feed Control Official (AAFCO).

MIXING DIRECTIONS

Thoroughly mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for turkey poults containing 400 grams chlortetracycline per ton.

The following table provides examples of mixing rates:

Type B CTC concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C CTC concentration (g/ton)
10,000 (5 g/lb)	80	1,920	400
40,000 (20 g/lb)	20	1,980	400
80,000 (40 g/lb)	10	1,990	400

The resulting Type C medicated feed should be fed continuously as the sole ration to turkey poults not over 4 weeks of age.

WITHDRAWAL PERIODS AND RESIDUE WARNINGS

No withdrawal period is required when used according to labeling.
Do not feed to turkeys producing eggs for human consumption.

USER SAFETY WARNINGS

Not for use in humans. Keep out of reach of children.

QUESTIONS/COMMENTS?

For technical assistance or to obtain a Safety Data Sheet(s), contact Pharmgate Animal Health LLC at 1-800-380-6099 or www.pharmgate.com. To report side effects, contact Pharmgate Animal Health LLC at 1-833-531-0114. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or http://www.fda.gov/reportanimalae.

Approved by FDA under ANADA # 200-510

Lot Number: _____

MANUFACTURED BY

Blue Bird Feed Mill
City, State, Zip

NET WEIGHT ON BAG OR BULK

*The final printed medicated feed label must state a single drug concentration.

[a] Sourced from Deracin™ (chlortetracycline Type A medicated article), ANADA # 200-510.

Pharmgate Inc. [29 July 2025]