

Product Profile

Circo/MycoGard[®]

Porcine Circovirus Vaccine, Type 2, Killed Baculovirus Vector, *Mycoplasma hyopneumoniae* Bacterin

Product Description

- One-dose combination of porcine circovirus (PCV) Type 2b vaccine and *Mycoplasma hyopneumoniae* bacterin for vaccination of swine
- Inactivated trivalent-adjuvanted formulation using StimGard Plus[™] adjuvant
- Each vial contains 50 or 250 doses (1 mL/dose)



Advantages

- Ready to use (RTU) - no mixing required.
- Labor-saving 1-mL dose – easy on nursing piglets and staff.
- Control 2 diseases in 1 injection.
- Demonstrated heterologous circovirus protection.
- 3 different classes of adjuvants in StimGard Plus[™] help provide for an optimal immune response.

Dosage / Administration

- Shake well before use. Using aseptic technique, inject a single 1.0-mL dose intramuscularly (IM).

Indications / Uses

- For use in healthy piglets of all breeds, 10 days of age or older, against Porcine Circovirus, Type 2, and *Mycoplasma hyopneumoniae*.
- Visit USDA website for full label indications & technical information. (productdata.aphis.usda.gov).

Precautions

- CAUTION: For animal use only.
- Store at 2 to 8°C (35-45°F). Do not freeze. Shake well before use. Use entire contents upon opening.
- Do not vaccinate within 21 days before slaughter.
- Transient reaction may occur at the injection site. If anaphylaxis occurs, use epinephrine or equivalent. Contains merthiolate (Thimerosal) as a preservative.
- This product has not been tested in pregnant animals.

Packaging

- 50-mL or 250-mL vials.
- Available in custom sizes.

Ordering Information

- Prod. number:
25010-50 (50 mL)
25010-250 (250 mL)

Key Features

- Pig-proven under field conditions.
- StimGard Plus[™] trivalent adjuvant system helps optimize both cell-mediated and humoral immune responses.
- Safe; studies show a <0.3% reaction rate.
- Excellent syringeability.
- Advanced purification process allows low-volume injection for minimal tissue reaction.
- No mixing means less handling, mistakes and mess
- QR (quick response) codes on the label for inventory and tracking purposes.

The labeling contains complete use information, including any cautions and warnings. Always read, understand and follow the labeling and use directions. See the reverse side for use directions and additional information.



Pharmgate
ANIMAL HEALTH

1800 Sir Tyler Drive
Wilmington, NC | 28405
910.679.8364
Pharmgate.com

Circo/MycoGard[®]

Porcine Circovirus Vaccine, Type 2, Killed Baculovirus Vector, *Mycoplasma hyopneumoniae* Bacterin

Porcine Circovirus Vaccine,
Type 2, Killed Baculovirus
Vector, *Mycoplasma*
hyopneumoniae Bacterin

Circo/MycoGard[®] | 250 ML
250 DOSES



This product has been shown to be effective for the vaccination of healthy piglets, 10 days of age or older against Porcine Circovirus, Type 2, and *Mycoplasma hyopneumoniae*. To review efficacy and safety data, go to productdata.aphis.usda.gov.

ADMINISTRATION AND DOSAGE:

Dose: 1.0 ml intramuscularly (I.M.).

CAUTION: For animal use only. Do not mix with other products, except as specified on this label. In case of human exposure, contact a physician. Store at 2 to 8°C (35°F to 45°F). Do not freeze. Shake well before use. Use entire contents upon opening. Do not vaccinate within 21 days before slaughter. Transient reaction may occur at the injection site. If anaphylaxis occurs, use epinephrine or equivalent. Contains merthiolate (Thimerosal) as a preservative. This product has not been tested in pregnant animals. Revaccination interval not established.

Product No:

Serial No:

Exp. Date:

Manufactured by:

Pharmgate Biologics
2575 University Avenue
West, Suite 100,
St. Paul, MN 55114
612-256-0930

VLN 329 / PCN 49K5.R1

Visit USDA website for full label indications & technical information: productdata.aphis.usda.gov

Pharmgate Animal Health swine vaccines/bacterins.

Product	Doses	Pharmgate Product Number
MycoGard [®] (2 dose)	50	20020-50
MycoGard [®] (2 dose)	250	20020-250
MycoGard-1 [™] (1 dose)	50	21010-50
MycoGard-1 [™] (1 dose)	250	21010-250
CircoGard [®] (1 dose)	50	22010-50
CircoGard [®] (1 dose)	250	22010-250
Circo/MycoGard [®] (1 dose)	50	25010-50
Circo/MycoGard [®] (1 dose)	250	25010-250